

# Spaces with Ricci curvature bounded from below

Nicola Gigli

March 10, 2014

# Lessons

- ▶ Basics of optimal transport
- ▶ Definition of spaces with Ricci curvature bounded from below
- ▶ Analysis on spaces with Ricci curvature bounded from below

# Lessons

- ▶ Basics of optimal transport
- ▶ Definition of spaces with Ricci curvature bounded from below
- ▶ Analysis on spaces with Ricci curvature bounded from below

# On the definition of spaces with Ricci curvature bounded from below

- ▶ Introduction
- ▶ The gradient flow of the relative entropy w.r.t.  $W_2$
- ▶ The gradient flow of the Dirichlet energy w.r.t.  $L^2$
- ▶ The heat flow as gradient flow

# On the definition of spaces with Ricci curvature bounded from below

- ▶ Introduction
- ▶ The gradient flow of the relative entropy w.r.t.  $W_2$
- ▶ The gradient flow of the Dirichlet energy w.r.t.  $L^2$
- ▶ The heat flow as gradient flow

# Completion / Compactification

A common practice in various fields of mathematic is to start studying a certain class of 'smooth' or 'nice' objects, and to close it w.r.t. some relevant topology.

In general, the study of the limit objects turns out to be useful to understand the properties of the original ones.

## Gromov's plan

When the original class of objects are Riemannian manifolds with some curvature bounds, this program has been proposed by Gromov.

# Gromov's plan

When the original class of objects are Riemannian manifolds with some curvature bounds, this program has been proposed by Gromov.

	Topology	Synthetic notion
Bounds from above/below on <a href="#">sectional</a> curvature		
Bounds from below on the <a href="#">Ricci</a> curvature		

# Gromov's plan

When the original class of objects are Riemannian manifolds with some curvature bounds, this program has been proposed by Gromov.

	Topology	Synthetic notion
Bounds from above/below on <a href="#">sectional</a> curvature	Gromov-Hausdorff convergence	
Bounds from below on the <a href="#">Ricci</a> curvature		

# Gromov's plan

When the original class of objects are Riemannian manifolds with some curvature bounds, this program has been proposed by Gromov.

	Topology	Synthetic notion
Bounds from above/below on <a href="#">sectional</a> curvature	Gromov-Hausdorff convergence	Alexandrov spaces
Bounds from below on the <a href="#">Ricci</a> curvature		

## Gromov's plan

When the original class of objects are Riemannian manifolds with some curvature bounds, this program has been proposed by Gromov.

	Topology	Synthetic notion
Bounds from above/below on <a href="#">sectional</a> curvature	Gromov-Hausdorff convergence	Alexandrov spaces
Bounds from below on the <a href="#">Ricci</a> curvature	measured Gromov-Hausdorff convergence	

## Gromov's plan

When the original class of objects are Riemannian manifolds with some curvature bounds, this program has been proposed by Gromov.

	Topology	Synthetic notion
Bounds from above/below on <a href="#">sectional</a> curvature	Gromov-Hausdorff convergence	Alexandrov spaces
Bounds from below on the <a href="#">Ricci</a> curvature	measured Gromov-Hausdorff convergence	$CD(K, N)$ spaces / $RCD(K, N)$ spaces

# Aim of the game

- (1) To understand what it means for a metric measure space to have Ricci curvature bounded from below
- (2) To prove in the non-smooth setting 'all' the theorems valid for manifolds with  $\text{Ric} \geq K$ ,  $\dim \leq N$  and their limits
- (3) To better understand the geometry of smooth manifolds via the study of non-smooth objects

# The curvature condition

**Theorem (Sturm-VonRenesse '05)** - see also Otto-Villani and Cordero  
Erausquin-McCann-Schmuckenschlager

Let  $M$  be a smooth Riemannian manifold. Then the following are equivalent:

- i) The Ricci curvature of  $M$  is uniformly bounded from below by  $K$
- ii) The relative entropy functional is  $K$ -convex on the space  $(\mathcal{P}_2(M), W_2)$

# The curvature condition

**Theorem** (Sturm-VonRenesse '05) - see also Otto-Villani and Cordero Erasquin-McCann-Schmuckenschlager

Let  $M$  be a smooth Riemannian manifold. Then the following are equivalent:

- i) The Ricci curvature of  $M$  is uniformly bounded from below by  $K$
- ii) The relative entropy functional is  $K$ -convex on the space  $(\mathcal{P}_2(M), W_2)$

**Definition** (Lott-Villani and Sturm '06)  $(X, d, \mathfrak{m})$  has Ricci curvature bounded from below by  $K$  if the relative entropy is  $K$ -convex on  $(\mathcal{P}_2(X), W_2)$ . Called  $CD(K, \infty)$  spaces, in short.

# The curvature condition

**Theorem (Sturm-VonRenesse '05)** - see also Otto-Villani and Cordero Erasmus-McCann-Schmuckenschlager

Let  $M$  be a smooth Riemannian manifold. Then the following are equivalent:

- i) The Ricci curvature of  $M$  is uniformly bounded from below by  $K$
- ii) The relative entropy functional is  $K$ -convex on the space  $(\mathcal{P}_2(M), W_2)$

**Definition (Lott-Villani and Sturm '06)**  $(X, d, \mathfrak{m})$  has Ricci curvature bounded from below by  $K$  if the relative entropy is  $K$ -convex on  $(\mathcal{P}_2(X), W_2)$ . Called  $CD(K, \infty)$  spaces, in short.

Basic features of the  $CD(K, \infty)$  condition:

- ▶ Compatibility with the Riemannian case
- ▶ Stability w.r.t. mGH convergence
- ▶ More general  $CD(K, N)$  spaces can be introduced

## Finsler structures are included

Cordero-Erausquin, Villani, Sturm proved that  $(\mathbb{R}^d, \|\cdot\|, \mathcal{L}^d)$  is a  $CD(0, \infty)$  space (in fact  $CD(0, d)$ ) for any norm.

# Finsler structures are included

Cordero-Erausquin, Villani, Sturm proved that  $(\mathbb{R}^d, \|\cdot\|, \mathcal{L}^d)$  is a  $CD(0, \infty)$  space (in fact  $CD(0, d)$ ) for any norm.

Some differences between the Finsler and Riemannian worlds:

Analysis:

---

Tangent / cotangent spaces  
can't be identified

No natural Dirichlet form

Geometry:

---

no Abresch-Gromoll inequality

no Splitting theorem

# Some observations

- ▶ For a given Finsler manifold the following are equivalent:
  - ▶ The manifold is Riemannian
  - ▶ The Sobolev space  $W^{1,2}$  is Hilbert
  - ▶ The heat flow is linear
  
- ▶ The heat flow can be seen as:
  - ▶ Gradient flow of the Dirichlet energy w.r.t.  $L^2$
  - ▶ Gradient flow of the relative entropy w.r.t.  $W_2$

## The idea

Restrict to the class of  $CD(K, \infty)$  spaces with linear heat flow.

# The idea

Restrict to the class of  $CD(K, \infty)$  spaces with linear heat flow.

What we have to do to show this makes sense:

- ▶ understand what is the heat flow on  $CD(K, \infty)$  spaces
- ▶ show that such flow is stable w.r.t. mGH convergence.

# The idea

Restrict to the class of  $CD(K, \infty)$  spaces with linear heat flow.

What we have to do to show this makes sense:

- ▶ understand what is the heat flow on  $CD(K, \infty)$  spaces
- ▶ show that such flow is stable w.r.t. mGH convergence.

Plan pursued in:

G. '09

G., Kuwada, Ohta '10

Ambrosio, G., Savaré '11

# On the definition of spaces with Ricci curvature bounded from below

- ▶ Introduction
- ▶ The gradient flow of the relative entropy w.r.t.  $W_2$
- ▶ The gradient flow of the Dirichlet energy w.r.t.  $L^2$
- ▶ The heat flow as gradient flow

## Definition of Gradient Flow: the smooth case

Let  $(x_t) \subset \mathbb{R}^d$  a smooth curve and  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  a smooth functional.  
Then

$$\begin{aligned} f(x_0) - f(x_t) &\leq \int_0^t |x'_s| |\nabla f|(x_s) \, ds \\ &\leq \frac{1}{2} \int_0^t |x'_s|^2 + |\nabla f|^2(x_s) \, ds. \end{aligned}$$

## Definition of Gradient Flow: the smooth case

Let  $(x_t) \subset \mathbb{R}^d$  a smooth curve and  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  a smooth functional.  
Then

$$\begin{aligned} f(x_0) - f(x_t) &\leq \int_0^t |x'_s| |\nabla f|(x_s) \, ds \\ &\leq \frac{1}{2} \int_0^t |x'_s|^2 + |\nabla f|^2(x_s) \, ds. \end{aligned}$$

Therefore

$$\begin{aligned} x'_t &= -\nabla f(x_t), \quad \forall t \geq 0 \\ &\Downarrow \\ f(x_0) &= f(x_t) + \frac{1}{2} \int_0^t |x'_s|^2 + |\nabla f|^2(x_s) \, ds, \quad \forall t > 0. \end{aligned}$$

## Definition of Gradient Flow: the metric setting

- ▶  $|\dot{x}_t| := \lim_{h \rightarrow 0} \frac{d(x_{t+h}, x_t)}{|h|}$  for an abs.cont. curve  $(x_t)$
- ▶  $|\partial^- F|(x) := \overline{\lim}_{y \rightarrow x} \frac{(F(x) - F(y))^+}{d(x, y)}$
- ▶ The weak chain rule

$$F(x_0) \leq F(x_t) + \frac{1}{2} \int_0^t |\dot{x}_s|^2 + |\partial^- F|^2(x_s) \, ds, \quad \forall t > 0.$$

holds for  $K$ -convex and l.s.c.  $F : X \rightarrow \mathbb{R} \cup \{+\infty\}$ .

## Definition of Gradient Flow: the metric setting

- ▶  $|\dot{x}_t| := \lim_{h \rightarrow 0} \frac{d(x_{t+h}, x_t)}{|h|}$  for an abs.cont. curve  $(x_t)$
- ▶  $|\partial^- F|(x) := \overline{\lim}_{y \rightarrow x} \frac{(F(x) - F(y))^+}{d(x, y)}$
- ▶ The weak chain rule

$$F(x_0) \leq F(x_t) + \frac{1}{2} \int_0^t |\dot{x}_s|^2 + |\partial^- F|^2(x_s) ds, \quad \forall t > 0.$$

holds for  $K$ -convex and l.s.c.  $F : X \rightarrow \mathbb{R} \cup \{+\infty\}$ .

**Definition**  $(x_t)$  is a Gradient Flow for the  $K$ -conv. and l.s.c. functional  $F$  provided  $(x_t) \subset \{F < \infty\}$  is a loc.abs.cont. curve and

$$F(x_0) = F(x_t) + \frac{1}{2} \int_0^t |\dot{x}_s|^2 + |\partial^- F|^2(x_s) ds, \quad \forall t > 0.$$

# General results about GF of $K$ -convex functionals

**Existence** Granted if the space is compact and  $F(x_0) < \infty$  (Ambrosio, G., Savaré '04 (after De Giorgi))

**Uniqueness** False in general

# Basic facts about the GF of the Entropy

**Thm. (G. '09)**

Let  $(X, d, \mathfrak{m})$  be a compact  $CD(K, \infty)$  space.

Then for  $\mu \in \mathcal{P}_2(X)$  with  $\text{Ent}_{\mathfrak{m}}(\mu) < \infty$  the GF of  $\text{Ent}_{\mathfrak{m}}$  starting from  $\mu$  exists and is unique.

# Basic facts about the GF of the Entropy

## Thm. (G. '09)

Let  $(X, d, \mathfrak{m})$  be a compact  $CD(K, \infty)$  space.

Then for  $\mu \in \mathcal{P}_2(X)$  with  $\text{Ent}_{\mathfrak{m}}(\mu) < \infty$  the GF of  $\text{Ent}_{\mathfrak{m}}$  starting from  $\mu$  exists and is unique.

Furthermore, such flow is stable w.r.t. mGH-convergence of the base space.

## mGH convergence of compact spaces

$(X_n, d_n, \mathbf{m}_n)$  converges to  $(X_\infty, d_\infty, \mathbf{m}_\infty)$  in the mGH sense if there is  $(Y, d_Y)$  and isometric embeddings  $\iota_n, \iota_\infty$  of the  $X$ 's into  $Y$  such that

$$(\iota_n)_\# \mathbf{m}_n \quad \text{weakly converges to} \quad (\iota_\infty)_\# \mathbf{m}_\infty$$

## mGH convergence of compact spaces

$(X_n, d_n, \mathbf{m}_n)$  converges to  $(X_\infty, d_\infty, \mathbf{m}_\infty)$  in the mGH sense if there is  $(Y, d_Y)$  and isometric embeddings  $\iota_n, \iota_\infty$  of the  $X$ 's into  $Y$  such that

$$(\iota_n)_\# \mathbf{m}_n \quad \text{weakly converges to} \quad (\iota_\infty)_\# \mathbf{m}_\infty$$

We say that  $n \mapsto \mu_n \in \mathcal{P}(X_n)$  weakly converges to  $\mu_\infty \in \mathcal{P}(X_\infty)$  provided

$$(\iota_n)_\# \mu_n \quad \text{weakly converges to} \quad (\iota_\infty)_\# \mu_\infty$$

# $\Gamma$ -convergence of the entropies

## Thm. (Lott-Sturm-Villani)

Let  $(X_n, d_n, \mathbf{m}_n)$  be converging to  $(X_\infty, d_\infty, \mathbf{m}_\infty)$ . Then:

- ▶  $\Gamma$  –  $\underline{\lim}$  inequality: for every sequence  $n \mapsto \mu_n \in \mathcal{P}(X_n)$  weakly converging to  $\mu_\infty \in \mathcal{P}(X_\infty)$  we have

$$\text{Ent}_{\mathbf{m}_\infty}(\mu_\infty) \leq \underline{\lim}_{n \rightarrow \infty} \text{Ent}_{\mathbf{m}_n}(\mu_n).$$

- ▶  $\Gamma$  –  $\overline{\lim}$  inequality: for every  $\mu_\infty \in \mathcal{P}(X_\infty)$  there is a sequence  $n \mapsto \mu_n \in \mathcal{P}(X_n)$  weakly converging to  $\mu_\infty$  such that

$$\text{Ent}_{\mathbf{m}_\infty}(\mu_\infty) \geq \overline{\lim}_{n \rightarrow \infty} \text{Ent}_{\mathbf{m}_n}(\mu_n).$$

# $\Gamma$ -convergence of the entropies

## Thm. (Lott-Sturm-Villani)

Let  $(X_n, d_n, \mathbf{m}_n)$  be converging to  $(X_\infty, d_\infty, \mathbf{m}_\infty)$ . Then:

- ▶  $\Gamma$  – lim inequality: for every sequence  $n \mapsto \mu_n \in \mathcal{P}(X_n)$  weakly converging to  $\mu_\infty \in \mathcal{P}(X_\infty)$  we have

$$\text{Ent}_{\mathbf{m}_\infty}(\mu_\infty) \leq \underline{\lim}_{n \rightarrow \infty} \text{Ent}_{\mathbf{m}_n}(\mu_n).$$

- ▶  $\Gamma$  –  $\overline{\text{lim}}$  inequality: for every  $\mu_\infty \in \mathcal{P}(X_\infty)$  there is a sequence  $n \mapsto \mu_n \in \mathcal{P}(X_n)$  weakly converging to  $\mu_\infty$  such that

$$\text{Ent}_{\mathbf{m}_\infty}(\mu_\infty) \geq \overline{\lim}_{n \rightarrow \infty} \text{Ent}_{\mathbf{m}_n}(\mu_n).$$

**Cor.** The  $CD(K, \infty)$  condition is closed w.r.t. mGH convergence.

## $\Gamma$ – lim for the slopes

**Thm. (G. '09)** Let  $X_n$  be  $CD(K, \infty)$  spaces mGH-converging to  $X_\infty$  and  $\mu_n$  weakly converging to  $\mu_\infty$ . Then

$$|\partial^- \text{Ent}_{\mathbf{m}_\infty}|(\mu_\infty) \leq \varinjlim_{n \rightarrow \infty} |\partial^- \text{Ent}_{\mathbf{m}_n}|(\mu_n).$$

## $\Gamma$ – lim for the slopes

**Thm. (G. '09)** Let  $X_n$  be  $CD(K, \infty)$  spaces mGH-converging to  $X_\infty$  and  $\mu_n$  weakly converging to  $\mu_\infty$ . Then

$$|\partial^- \text{Ent}_{\mathbf{m}_\infty}|(\mu_\infty) \leq \underline{\lim}_{n \rightarrow \infty} |\partial^- \text{Ent}_{\mathbf{m}_n}|(\mu_n).$$

**Cor. 1** Let  $X_n$  be mGH-converging to  $X_\infty$  and  $\mu_n$  be weakly converging to  $\mu_\infty$  be such that

$$\lim_{n \rightarrow \infty} \text{Ent}_{\mathbf{m}_n}(\mu_n) = \text{Ent}_{\mathbf{m}_\infty}(\mu_\infty) < \infty.$$

Then the GF of  $\text{Ent}_{\mathbf{m}_n}$  starting from  $\mu_n$  converge to the GF of  $\text{Ent}_{\mathbf{m}_\infty}$  starting from  $\mu_\infty$ .

## $\Gamma$ – lim for the slopes

**Thm. (G. '09)** Let  $X_n$  be  $CD(K, \infty)$  spaces mGH-converging to  $X_\infty$  and  $\mu_n$  weakly converging to  $\mu_\infty$ . Then

$$|\partial^- \text{Ent}_{\mathbf{m}_\infty}|(\mu_\infty) \leq \underline{\lim}_{n \rightarrow \infty} |\partial^- \text{Ent}_{\mathbf{m}_n}|(\mu_n).$$

**Cor. 1** Let  $X_n$  be mGH-converging to  $X_\infty$  and  $\mu_n$  be weakly converging to  $\mu_\infty$  be such that

$$\lim_{n \rightarrow \infty} \text{Ent}_{\mathbf{m}_n}(\mu_n) = \text{Ent}_{\mathbf{m}_\infty}(\mu_\infty) < \infty.$$

Then the GF of  $\text{Ent}_{\mathbf{m}_n}$  starting from  $\mu_n$  converge to the GF of  $\text{Ent}_{\mathbf{m}_\infty}$  starting from  $\mu_\infty$ .

**Cor. 2** The condition ' $CD(K, \infty)$ +linearity of the GF of the entropy' is closed w.r.t. mGH convergence.

# On the definition of spaces with Ricci curvature bounded from below

- ▶ Introduction
- ▶ The gradient flow of the relative entropy w.r.t.  $W_2$
- ▶ The gradient flow of the Dirichlet energy w.r.t.  $L^2$
- ▶ The heat flow as gradient flow

## Variational definition of $|Df|$ on $\mathbb{R}^d$

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be smooth.

Then  $|Df|$  is the minimum continuous function  $G$  for which

$$|f(\gamma_1) - f(\gamma_0)| \leq \int_0^1 G(\gamma_t) |\dot{\gamma}_t| dt$$

holds for any smooth curve  $\gamma$

# Test plans

Let  $\pi \in \mathcal{P}(C([0, 1], X))$ . We say that  $\pi$  is a test plan provided:

- ▶ for some  $C > 0$  it holds

$$e_{t\#}\pi \leq C\mathbf{m}, \quad \forall t \in [0, 1].$$

- ▶ it holds

$$\iint_0^1 |\dot{\gamma}_t|^2 dt d\pi < \infty$$

## The Sobolev class $S^2(X, d, \mathbf{m})$

We say that  $f : X \rightarrow \mathbb{R}$  belongs to  $S^2(X, d, \mathbf{m})$  provided there exists  $G \in L^2(X, \mathbf{m})$ ,  $G \geq 0$  such that

$$\int |f(\gamma_1) - f(\gamma_0)| \, d\pi(\gamma) \leq \iint_0^1 G(\gamma_t) |\dot{\gamma}_t| \, dt \, d\pi(\gamma)$$

for any test plan  $\pi$ .

Any such  $G$  is called weak upper gradient of  $f$ .

The minimal  $G$  in the  $\mathbf{m}$ -a.e. sense will be denoted by  $|Df|$

# Basic properties

**Lower semicontinuity** From  $f_n \rightarrow f$  **m-a.e.** with  $f_n \in S^2$  and  $|Df_n| \rightarrow G$  weakly in  $L^2$  we deduce

$$f \in S^2, \quad |Df| \leq G$$

**Locality**

$$|Df| = |Dg| \quad \mathbf{m-a.e. on} \{f = g\}$$

**Chain rule**

$$|D(\varphi \circ f)| = |\varphi'| \circ f |Df|$$

for  $\varphi$  Lipschitz

**'Leibniz rule'**

$$|D(fg)| \leq |f| |Dg| + |g| |Df|$$

for  $f, g \in S^2 \cap L^\infty$

# The Energy $E$ and the Sobolev space $W^{1,2}$

We define  $E : L^2(X, \mathbf{m}) \rightarrow [0, +\infty]$  as

$$E(f) := \begin{cases} \frac{1}{2} \int |Df|^2 \, d\mathbf{m} & \text{if } f \in \mathcal{S}^2(X), \\ +\infty & \text{otherwise.} \end{cases}$$

Then  $E$  is convex and lower semicontinuous.

# The Energy $E$ and the Sobolev space $W^{1,2}$

We define  $E : L^2(X, \mathbf{m}) \rightarrow [0, +\infty]$  as

$$E(f) := \frac{1}{2} \int |Df|^2 d\mathbf{m} \quad \text{if } f \in \mathcal{S}^2(X), \quad +\infty \quad \text{otherwise.}$$

Then  $E$  is convex and lower semicontinuous.

The Sobolev space  $W^{1,2}(X)$  is  $W^{1,2}(X) := L^2(X) \cap \mathcal{S}^2(X)$  endowed with the norm

$$\|f\|_{W^{1,2}}^2 := \|f\|_{L^2}^2 + \|Df\|_{L^2}^2$$

$W^{1,2}(X)$  is a Banach space.

## Laplacian (first definition)

We say that  $f \in D(\Delta) \subset W^{1,2}(X)$  if  $\partial^- E(f) \neq 0$ .

In this case we define  $\Delta f := -v$ , where  $v$  is the element of minimal norm in  $\partial^- E(f)$ .

## 'Integration by parts'

For  $f \in D(\Delta)$  and  $g \in W^{1,2}(X)$  we have

$$\left| \int g \Delta f \, d\mathbf{m} \right| \leq \int |Dg| |Df| \, d\mathbf{m}.$$

For a  $C^1$  map  $u : \mathbb{R} \rightarrow \mathbb{R}$  we have

$$\int u(f) \Delta f \, d\mathbf{m} = - \int u'(f) |Df|^2 \, d\mathbf{m}.$$

## Gradient flow of $E$ w.r.t. $L^2$

For any  $f_0 \in L^2(X, \mathfrak{m})$  there exists a unique map  $t \mapsto f_t \in L^2(X, \mathfrak{m})$  such that

$$\frac{d^+}{dt} f_t = \Delta f_t, \quad \forall t > 0.$$

# On the definition of spaces with Ricci curvature bounded from below

- ▶ Introduction
- ▶ The gradient flow of the relative entropy w.r.t.  $W_2$
- ▶ The gradient flow of the Dirichlet energy w.r.t.  $L^2$
- ▶ The heat flow as gradient flow

# The result

**Thm.** (G., Kuwada, Ohta '10. Ambrosio, G. Savaré '11)

Let  $(X, d, \mathbf{m})$  be a  $CD(K, \infty)$  space and  $\mu = f\mathbf{m} \in \mathcal{P}_2(X)$  with  $f \in L^2(X, \mathbf{m})$ . Let

- ▶  $t \mapsto f_t$  be the GF of  $E$  w.r.t.  $L^2$  starting from  $f$
- ▶  $t \mapsto \mu_t$  be the GF of  $\text{Ent}_{\mathbf{m}}$  w.r.t.  $W_2$  starting from  $\mu$

Then

$$\mu_t = f_t \mathbf{m} \quad \forall t \geq 0.$$

# The result

**Thm.** (G., Kuwada, Ohta '10. Ambrosio, G. Savaré '11)

Let  $(X, d, \mathbf{m})$  be a  $CD(K, \infty)$  space and  $\mu = f\mathbf{m} \in \mathcal{P}_2(X)$  with  $f \in L^2(X, \mathbf{m})$ . Let

- ▶  $t \mapsto f_t$  be the GF of  $E$  w.r.t.  $L^2$  starting from  $f$
- ▶  $t \mapsto \mu_t$  be the GF of  $\text{Ent}_{\mathbf{m}}$  w.r.t.  $W_2$  starting from  $\mu$

Then

$$\mu_t = f_t \mathbf{m} \quad \forall t \geq 0.$$

Idea of the proof: to show that  $t \mapsto f_t \mathbf{m}$  is a GF of  $\text{Ent}_{\mathbf{m}}$  w.r.t.  $W_2$  and conclude by uniqueness of the latter.

# Spaces with Riemannian Ricci curvature bounded from below

$$\begin{aligned} RCD(K, N) &:= CD(K, N) + \text{linearity of the heat flow} \\ &= CD(K, N) + W^{1,2} \text{ is Hilbert} \end{aligned}$$

Thank you

# Basic properties of the GF of $E$

**Mass preservation**  $\int f_t \, d\mathbf{m} = \int f_0 \, d\mathbf{m}$  for every  $t \geq 0$

**Weak maximum principle** If  $f_0 \leq C$  then  $f_t \leq C$  for every  $t \geq 0$

**Entropy dissipation** For  $0 < c \leq f_0 \leq C$  the map  $t \mapsto \int f_t \log f_t \, d\mathbf{m}$  is absolutely continuous and it holds

$$\frac{d}{dt} \int f_t \log f_t \, d\mathbf{m} = - \int \frac{|Df_t|^2}{f_t} \, d\mathbf{m}$$

# A non-trivial property of the heat flow - Kuwada's lemma

Suppose that  $\mu_0 := f\mathbf{m}$  is in  $\mathcal{P}_2(X)$ .

# A non-trivial property of the heat flow - Kuwada's lemma

Suppose that  $\mu_0 := f \mathbf{m}$  is in  $\mathcal{P}_2(X)$ .

Then  $\mu_t := f_t \mathbf{m}$  is in  $\mathcal{P}_2(X)$  and the curve  $t \mapsto \mu_t$  is absolutely continuous w.r.t.  $W_2$  and

$$|\dot{\mu}_t|^2 \leq \int \frac{|Df_t|^2}{f_t} \, d\mathbf{m}$$