

EURANDOM PREPRINT SERIES

2016-014

November 16, 2016

Metastability on the Hierarchical Lattice

F. den Hollander, O. Jovanovski
ISSN 1389-2355

METASTABILITY ON THE HIERARCHICAL LATTICE

FRANK DEN HOLLANDER, OLIVER JOVANOVSKI

ABSTRACT. We study metastability for Glauber spin-flip dynamics on the N -dimensional hierarchical lattice with n hierarchical levels. Each vertex carries an Ising spin that can take the values -1 or $+1$. Spins interact with an external magnetic field $h > 0$. Pairs of spins interact with each other according to a ferromagnetic pair potential $\vec{J} = \{J_i\}_{i=1}^n$, where $J_i > 0$ is the strength of the interaction between spins at hierarchical distance i . Spins flip according to a Metropolis dynamics at inverse temperature β . In the limit as $\beta \rightarrow \infty$, we analyse the crossover time from the metastable state \boxminus (all spins -1) to the stable state \boxplus (all spins $+1$). Under the assumption that \vec{J} is non-increasing, we identify the mean transition time up to a multiplicative factor $1 + o_\beta(1)$. On the scale of its mean, the transition time is exponentially distributed. We also identify the set of configurations representing the gate for the transition. For the special case where $J_i = \tilde{J}/N^i$, $1 \leq i \leq n$, with $\tilde{J} > 0$ the relevant formulas simplify considerably. Also the hierarchical mean-field limit $N \rightarrow \infty$ can be analysed in detail.

1. INTRODUCTION

Interacting particle systems evolving according to a *Metropolis dynamics* associated with an energy functional, called the *Hamiltonian*, may end up being trapped for a long time near a state that is a local minimum but not a global minimum. Just how long it takes for the system to escape from the energy valley around a local minimum and reach the global minimum depends on how deep this valley is. The deepest local minima are called *metastable states*, the global minimum is called the *stable state*. While being trapped near a metastable state, the system is said to be in a quasi-equilibrium. The transition to the stable state marks the relaxation of the system to equilibrium. To describe this relaxation, it is of interest to compute the transition time and to identify the set of critical configurations the system has to cross in order to achieve the transition. The critical configurations constitute the lowest saddle points in the energy landscape encountered along paths that achieve the crossover.

Metastability for interacting particle systems on *lattices* has been studied intensively in the past three decades. Various different approaches have been proposed, which are summarised in the monographs by Olivieri and Vares ^{ov05} [5], Bovier and den Hollander ^{dh15} [1]. Recently, there has been interest in metastability for interacting particle systems on *random graphs*, which is much more challenging because the transition time depends in a delicate manner on the realisation of the graph.

In the present paper we are interested in metastability for Glauber spin-flip dynamics on the N -dimensional *hierarchical lattice* at low temperature. We obtain a full description of both the transition time and the set of critical configurations representing the gate for the transition. Our results are part of a larger enterprise in which the goal is to understand metastability on large graphs. The hierarchical lattice is interesting because it has a non-trivial geometric structure and allows for a rich variability in the choice of the interaction parameters.

The outline of the paper is as follows. In Section 1.1 we recall the definition of Glauber spin-flip dynamics on an arbitrary finite connected graph. In Section 1.2 we recall the basic geometric definitions that are needed for the description of metastability and recall three key theorems from the literature that are valid in the limit of low temperature. These theorems, which are based on a certain *key hypothesis* but are otherwise model-independent, state that the mean transition time equals $[1 + o_\beta(1)] K^* e^{\beta\Gamma^*}$, with β the inverse temperature, and that the gate for the transition

Date: November 16, 2016.

is \mathcal{C}^* , where $(\Gamma^*, \mathcal{C}^*, K^*)$ is a model-dependent triple. The theorems also show that the spectral gap of the generator of the dynamics scales like the inverse of the mean transition time and that the transition time divided by its mean is exponentially distributed asymptotically. In Section 1.3 we recall that the prefactor K^* is given by a variational formula. In Section 1.4 we define the hierarchical lattice. In Section 1.5 we verify the key hypothesis for Glauber spin-flip dynamics on the hierarchical lattice and state *five assumptions* on the interaction parameters. In Section 1.6 we state our main theorems, which identify the triple $(\Gamma^*, \mathcal{C}^*, K^*)$ for the hierarchical lattice subject to these assumptions. In Section 1.7 we close with a discussion and point to related literature. The proofs of the main theorems are given in Sections 2–4. The framework that is recalled in Sections 1.1–1.3 is taken from Bovier and den Hollander ^{bdh15}[1, Chapter 16].

1.1. Ising model and Glauber spin-flip dynamics. Given a finite connected graph $G = (V, E)$, let $\Omega = \{-1, +1\}^V$ be the set of configurations $\sigma = \{\sigma(v) : v \in V\}$ that assigns to each vertex $v \in V$ a spin-value $\sigma(v) \in \{-1, +1\}$. Two configurations that will be of particular interest to us are those where all spins point up, respectively, down:

$$(1.1) \quad \boxplus \equiv +1, \quad \boxminus \equiv -1.$$

For $\beta \geq 0$, playing the role of *inverse temperature*, we define the Gibbs measure

$$(1.2) \quad \mu_\beta(\sigma) = \frac{1}{Z_\beta} e^{-\beta \mathcal{H}(\sigma)}, \quad \sigma \in \Omega,$$

where $\mathcal{H} : \Omega \rightarrow \mathbb{R}$ is the *Hamiltonian* that assigns an energy to each configuration given by

$$(1.3) \quad \mathcal{H}(\sigma) = -\frac{1}{2} \sum_{(v,w) \in E} J_{(v,w)} \sigma(v)\sigma(w) - \frac{h}{2} \sum_{v \in V} \sigma(v), \quad \sigma \in \Omega,$$

where $\vec{J} = \{J_e\}_{e \in E}$ is the *ferromagnetic pair potential* acting along edges, satisfying $J_e \geq 0$ for all $e \in E$, and $h > 0$ is the *external magnetic field*.

For two configurations $\sigma, \eta \in \Omega$, we write $\sigma \sim \eta$ when σ and η agree at all but one vertex. A transition from σ to η corresponds to a flip of a single spin, and is referred to as an *allowed move*. Glauber spin-flip dynamics on Ω is the continuous-time Markov process $(\sigma_t)_{t \geq 0}$ defined by the transition rates

$$(1.4) \quad c_\beta(\sigma, \eta) = \begin{cases} e^{-\beta[\mathcal{H}(\eta) - \mathcal{H}(\sigma)]_+}, & \sigma \sim \eta, \\ 0, & \text{otherwise.} \end{cases}$$

The Gibbs measure in (1.2) is the *reversible equilibrium* of this dynamics. We write $P_\sigma^{G, \beta}$ to denote the law of $(\sigma_t)_{t \geq 0}$ given $\sigma_0 = \sigma$, $\mathcal{L}^{G, \beta}$ to denote the associated generator, and $\lambda^{G, \beta}$ to denote the principal eigenvalue of $\mathcal{L}^{G, \beta}$. The upper indices G, β exhibit the dependence on the underlying graph G and the interaction strength β between neighbouring spins. For $A \subseteq \Omega$, we write

$$(1.5) \quad \tau_A = \inf \{t > 0 : \sigma_t \in A, \exists 0 < s < t : \sigma_s \neq \sigma_0\}$$

to denote the first hitting time of the set A after the starting configuration is left.

1.2. Metastability. To describe the metastable behaviour of our dynamics we need the following geometric definitions.

Definition 1.1. (a) *The communication height between two distinct configurations $\sigma, \eta \in \Omega$ is*

$$(1.6) \quad \Phi(\sigma, \eta) = \min_{\gamma: \sigma \rightarrow \eta} \max_{\xi \in \gamma} \mathcal{H}(\xi),$$

where the minimum is taken over all paths $\gamma : \sigma \rightarrow \eta$ consisting of allowed moves only. The communication height between two non-empty disjoint sets $A, B \subset \Omega$ is

$$(1.7) \quad \Phi(A, B) = \min_{\sigma \in A, \eta \in B} \Phi(\sigma, \eta).$$

(b) The stability level of $\sigma \in \Omega$ is

$$(1.8) \quad V_\sigma = \min_{\substack{\eta \in \Omega: \\ \mathcal{H}(\eta) < \mathcal{H}(\sigma)}} \Phi(\sigma, \eta) - \mathcal{H}(\sigma).$$

(c) The set of stable configurations is

$$(1.9) \quad \Omega_{\text{stab}} = \left\{ \sigma \in \Omega: \mathcal{H}(\sigma) = \min_{\eta \in \Omega} \mathcal{H}(\eta) \right\}.$$

(d) The set of metastable configurations is

$$(1.10) \quad \Omega_{\text{meta}} = \left\{ \sigma \in \Omega \setminus \Omega_{\text{stab}}: V_\sigma = \max_{\eta \in \Omega \setminus \Omega_{\text{stab}}} V_\eta \right\}.$$

It is easy to check that $\Omega_{\text{stab}} = \{\boxplus\}$ for all G because $h > 0$ and $J_e \geq 0$ for all $e \in E$. In general, Ω_{meta} is not a singleton. In order to proceed, we need the following *key hypothesis*:

$$(1.11) \quad \text{(H)} \quad \Omega_{\text{meta}} = \{\boxminus\}.$$

Hypothesis (H) states that $\{\boxminus, \boxplus\}$ is a metastable pair. The energy barrier between \boxminus and \boxplus is

$$(1.12) \quad \Gamma^* = \Phi(\boxminus, \boxplus) - \mathcal{H}(\boxminus),$$

which is a key quantity for the description of the metastable behaviour of our dynamics. We will say that a path $\gamma: \boxminus \rightarrow \boxplus$ is an *optimal path* when

$$(1.13) \quad \max_{\eta \in \gamma} \mathcal{H}(\eta) = \Phi(\boxminus, \boxplus).$$




FIGURE 1. Schematic picture of the protocritical set and the critical set.

Definition 1.2. Let $(\mathcal{P}^*, \mathcal{C}^*)$ be the unique maximal subset of $\Omega \times \Omega$ with the following properties (see Fig. 1):

- (1) $\forall \sigma \in \mathcal{P}^* \exists \eta \in \mathcal{C}^*: \sigma \sim \eta,$
 $\forall \eta \in \mathcal{C}^* \exists \sigma \in \mathcal{P}^*: \eta \sim \sigma.$
- (2) $\forall \sigma \in \mathcal{P}^*: \Phi(\sigma, \boxminus) < \Phi(\sigma, \boxplus).$
- (3) $\forall \sigma \in \mathcal{C}^* \exists \gamma: \sigma \rightarrow \boxplus:$
 - (i) $\max_{\eta \in \gamma} \mathcal{H}(\eta) \leq \Phi(\boxminus, \boxplus).$
 - (ii) $\gamma \cap \{\eta \in \Omega: \Phi(\eta, \boxminus) < \Phi(\eta, \boxplus)\} = \emptyset.$

Think of \mathcal{P}^* as the set of configurations where the dynamics, on its way from \boxminus to \boxplus , is ‘almost at the top’, and of \mathcal{C}^* as the set of configurations where it is ‘at the top and capable of crossing over’. We refer to \mathcal{P}^* as the *protocritical set* and to \mathcal{C}^* as the *critical set*. Uniqueness follows from the observation that if $(\mathcal{P}_1^*, \mathcal{C}_1^*)$ and $(\mathcal{P}_2^*, \mathcal{C}_2^*)$ both satisfy conditions (1)–(3), then so does $(\mathcal{P}_1^* \cup \mathcal{P}_2^*, \mathcal{C}_1^* \cup \mathcal{C}_2^*)$. Note that

$$(1.14) \quad \begin{aligned} \mathcal{H}(\sigma) < \Phi(\boxminus, \boxplus) & \quad \forall \sigma \in \mathcal{P}^*, \\ \mathcal{H}(\sigma) = \Phi(\boxminus, \boxplus) & \quad \forall \sigma \in \mathcal{C}^*. \end{aligned}$$

It is shown in Bovier and den Hollander [1, Chapter 16]^{BdH15} that *subject to hypothesis (H)* the following three theorems hold.

Theorem 1.3. $\lim_{\beta \rightarrow \infty} P_{\square}^{G,\beta}(\tau_{C^*} < \tau_{\square} \mid \tau_{\square} < \tau_{\boxplus}) = 1.$

Theorem 1.4. *There exists a $K^* \in (0, \infty)$ such that*

$$(1.15) \quad \lim_{\beta \rightarrow \infty} e^{-\beta \Gamma^*} E_{\square}^{G,\beta}(\tau_{\boxplus}) = K^*.$$

Theorem 1.5. (a) $\lim_{\beta \rightarrow \infty} \lambda^{G,\beta} E_{\square}^{G,\beta}(\tau_{\boxplus}) = 1.$

(b) $\lim_{\beta \rightarrow \infty} P_{\square}^{G,\beta}(\tau_{\boxplus}/E_{\square}^{G,\beta}(\tau_{\boxplus}) > t) = e^{-t}$ for all $t \geq 0.$




FIGURE 2. Schematic picture of \mathcal{H} , \square , \boxplus , Γ^* and C^* . Lemma 1.6 shows that $1/K^*$ is in essence proportional to $|C^*|$.

The proofs of Theorems 1.3–1.5 in [1]^{BdH15} do not rely on the details of the graph G , provided it is finite, connected and non-oriented. For concrete choices of G , the task is to verify hypothesis (H) and to identify the triple

$$(1.16) \quad (\Gamma^*, C^*, K^*).$$

A schematic picture of the role of these quantities is given in Fig. 2.

1.3. Variational formula for the prefactor. The prefactor K^* in Theorem 1.4 is given by a variational formula (see [1, Lemma 16.17]^{BdH15}):

$$(1.17) \quad \frac{1}{K^*} = \min_{C_1, \dots, C_I} \min_{f: S^* \rightarrow [0,1]; f|_{S_{\square}} \equiv 1, f|_{S_{\boxplus}} \equiv 0, f|_{S_k} = C_k} \frac{1}{2} \sum_{\sigma, \eta \in S^*} \mathbf{1}_{\{\sigma \sim \eta\}} [f(\sigma) - f(\eta)]^2.$$

Here, $\{S_k\}_{k=1}^I$ is the unique sequence of maximally connected disjoint sets $S_k \subseteq \Omega$ defined by

$$(1.18) \quad \sigma \in S_k \iff \mathcal{H}(\sigma) < \Phi(\square, \boxplus), \Phi(\sigma, \square) = \Phi(\sigma, \boxplus) = \Phi(\square, \boxplus).$$

Think of $\{S_k\}_{k=1}^I$ as ‘wells at the top’ (see Fig. 3). The sets $S_{\square}, S_{\boxplus}$ are defined by

$$(1.19) \quad \begin{aligned} S_{\square} &= \{\sigma \in \Omega: \Phi(\sigma, \square) < \Phi(\square, \boxplus)\}, \\ S_{\boxplus} &= \{\sigma \in \Omega: \Phi(\sigma, \boxplus) < \Phi(\square, \boxplus)\}, \end{aligned}$$

and are to be thought of as the ‘valleys’ around \square and \boxplus . The set S^* is defined by

$$(1.20) \quad S^* = \{\sigma \in \Omega: \Phi(\sigma, \square) \vee \Phi(\sigma, \boxplus) \leq \Phi(\square, \boxplus)\},$$

i.e., the maximally connected set with energy $\leq \Phi(\square, \boxplus)$ containing \square and \boxplus . Note that $\{S_k\}_{k=1}^I, S_{\square}, S_{\boxplus} \subseteq S^*$.

The variational problem in (1.17) has the interpretation of the *capacity* between S_{\square} and S_{\boxplus} for *simple random walk* on S^* jumping at rate 1 after the sets $\{S_k\}_{k=1}^I, S_{\square}, S_{\boxplus}$ are wired. If we impose *additional constraints* on the optimal paths and their behaviour near the set C^* , then (1.17) simplifies considerably, as is shown in the following lemma.




FIGURE 3. Schematic picture of the wells $\{S_k\}_{k=1}^I$. Note that $\mathcal{C}^* \subseteq S \setminus (S_\square \cup S_\boxplus)$.

Lemma 1.6. *Suppose that there exists a $k^* \in \mathbb{N}$ such that the following are true:*

- (i) $\mathcal{C}^* = \{\sigma \in S^* : |\sigma| = k^*\}$.
- (ii) For all $\sigma \in \mathcal{C}^*$ the sets

$$(1.21) \quad \begin{aligned} U_\sigma^- &= \{\eta \in S^* : \eta \sim \sigma, |\eta| = |\sigma| - 1\}, \\ U_\sigma^+ &= \{\eta \in S^* : \eta \sim \sigma, |\eta| = |\sigma| + 1\}, \end{aligned}$$

satisfy

eq:condition-lemma

$$(1.22) \quad \Phi(\eta, \square) < \Phi(\square, \boxplus) \quad \forall \eta \in U_\sigma^-, \quad \Phi(\eta, \boxplus) < \Phi(\square, \boxplus) \quad \forall \eta \in U_\sigma^+.$$

Then (1.17) simplifies to

eq:reduced-varform-0

$$(1.23) \quad \frac{1}{K^*} = \sum_{\sigma \in \mathcal{C}^*} \frac{|U_\sigma^-| |U_\sigma^+|}{|U_\sigma^-| + |U_\sigma^+|}.$$

Proof. The proof is analogous to that in [1, Section 17.5]. The variational problem in (1.17) simplifies because of the following two facts that are specific to Glauber dynamics:

- $S^* \setminus [S_\square \cup S_\boxplus] = \mathcal{C}^*$, i.e., there are no wells inside \mathcal{C}^* .
- There are no allowed moves within \mathcal{C}^* , i.e., critical configurations cannot transform into each other via single spin-flips.

Consequently, (1.17) reduces to

$$(1.24) \quad \frac{1}{K^*} = \min_{h: \mathcal{C}^* \rightarrow [0,1]} \sum_{\sigma \in \mathcal{C}^*} [1 - h(\sigma)] 2|U_\sigma^-| + [h(\sigma)] 2|U_\sigma^+|,$$

where U_σ^- and U_σ^+ consist of the configurations in S_\square and S_\boxplus , respectively, that can be reached from $\sigma \in \mathcal{C}^*$ by a single spin-flip. The solution of (1.24) is computed easily to obtain (1.23) \square

Remark 1.7. An immediate consequence of the additional assumptions in Lemma 1.6 is that $I = 0$ ('no wells at the top') and that all configurations in S^* that are neighbours of configurations in \mathcal{C}^* have an energy that is strictly below $\Phi(\square, \boxplus)$ ('the top is not flat'). Consequently, only transitions from \mathcal{C}^* to S_\square and S_\boxplus ('down from the top') contribute to the prefactor (see Fig. 4).




FIGURE 4. Configurations in \mathcal{C}^* are strict maxima in the energy profile of an optimal path. No plateau or wells are present.

1.4. The hierarchical lattice. Let $N \in \mathbb{N} \setminus \{1\}$, and define the N -dimensional hierarchical lattice Λ_N to be the metric space (\mathbb{N}, d) with \mathbb{N} the set of positive integers and d the ultrametric defined by

$$(1.25) \quad d(a, b) = \max \{k \in \mathbb{N}_0 : a \bmod N^k \neq b \bmod N^k\}, \quad a, b \in \mathbb{N},$$

which is called the *hierarchical distance*. We say that $A \subseteq \mathbb{N}$ is a k -block of Λ_N when $|A| = N^k$ and $d(a, b) \leq k$ for all $a, b \in A$. In particular, we define Λ_N^n to be the n -block

$$(1.26) \quad \Lambda_N^n = \{1, 2, \dots, N^n\},$$

which is the N -dimensional hierarchical lattice with n hierarchical levels (see Fig. 5).




FIGURE 5. Schematic representation of Λ_4^3 . The distance from the vertex in the lower-left corner to any vertex in the lower-left 1-block different from that vertex equals 1, to any vertex in the lower-left 2-block that is not in the lower-left 1-block equals 2, and to any vertex in the lower-left 3-block that is not in the lower-left 2-block equals 3. Note that, with this interpretation, for any two vertices v and w the size of the smallest box containing both v and w is $N^{d(v,w)}$.

The set Λ_N^n is the underlying graph from which we build our state space $\Omega = \{-1, +1\}^{\Lambda_N^n}$. We may alternatively write $\Lambda_N^n = \{v_1, \dots, v_{N^n}\}$ with v_a the vertex corresponding to the integer a . Note that $d(v_a, v_b) = d(a, b)$. We define $\gamma: \boxplus \rightarrow \boxtimes$ to be the path $\gamma = (\gamma_0, \dots, \gamma_{N^n})$, where γ_k is the configuration with $\gamma_k(v_a) = +1$ for $a \leq k$ and $\gamma_k(v_a) = -1$ for $a > k$, i.e., spins are flipped upward in the order in which they are labelled. We refer to γ as the *reference path*, and it will play a crucial role in our analysis.

Whenever convenient, we may think of Ω as the power set of Λ_N^n and of configurations $\sigma \in \Omega$ as subsets of Λ_N^n . Thus, we may identify a configuration $\sigma \in \{-1, +1\}^{\Lambda_N^n}$ with the set $\{v \in \Lambda_N^n : \sigma(v) = +1\}$ and its flipped image $\bar{\sigma}$ with the set $\{v \in \Lambda_N^n : \sigma(v) = -1\}$.

To define the interaction, we make Λ_N^n into a complete graph by placing an edge between all pairs $v, w \in \Lambda_N^n$ with $v \neq w$. The ferromagnetic pair potential between such pairs equals $J_{d(v,w)}$, where

$$(1.27) \quad \vec{J} = \{J_i\}_{i=1}^n$$

is chosen such that $J_i > 0$ for $1 \leq i \leq n$. Hence the Hamiltonian in (1.3) becomes

$$(1.28) \quad \mathcal{H}(\sigma) = -\frac{1}{2} \sum_{\substack{v, w \in \Lambda_N^n \\ v \neq w}} J_{d(v,w)} \sigma(v) \sigma(w) - \frac{h}{2} \sum_{v \in \Lambda_N^n} \sigma(v).$$

1.5. Hypothesis and Assumptions. We want to apply the theory behind Theorems 1.3–1.5, for which we need to verify Hypothesis (H) in (1.11). In the sequel we will need five assumptions on the interaction parameters of our model.

Assumption (A1):

$$(1.29) \quad \left(1 - \frac{1}{N}\right) \sum_{i=1}^n J_i N^i > h.$$

(A1) guarantees that Ξ is a local minimum and corresponds to the range of parameters for which the system is in the *metastable regime*.

Theorem 1.8. *Suppose that \vec{J} is monotone, i.e. either non-increasing or non-decreasing, and that (A1) holds. Then hypothesis (H) is verified.*

We will see from the proof of Theorem 1.8 that without (A1) there are no local minima in the energy landscape.

Our main task is to identify the triplet $(\Gamma^*, \mathcal{C}^*, K^*)$ in (1.16). To do so, we require *four assumptions* on \vec{J} , which we list below.

Assumption (A2):

$$(1.30) \quad \begin{aligned} \text{(a)} \quad & \exists \delta > 0, M \in \mathbb{N}: \quad 1 - \delta \geq [\hat{s}] - \hat{s} \geq \delta \quad \forall N \geq M, \\ \text{(b)} \quad & \liminf_{N \rightarrow \infty} \left| \sum_{i=\hat{m}+1}^n J_i N^i - h \right| > 0, \end{aligned}$$

where

$$(1.31) \quad \hat{m} = \max \left\{ 0 \leq m \leq n-1: \left(1 - \frac{1}{N}\right) \sum_{i=m+1}^n J_i N^i > h \right\},$$

$$(1.32) \quad \hat{s} = \frac{N}{2} (J_{\hat{m}+1} N^{\hat{m}+1})^{-1} \left[\left(1 - \frac{1}{N}\right) \sum_{i=\hat{m}+1}^n J_i N^i - h \right].$$

(A2)(a) guarantees that \hat{s} is not an integer when N is sufficiently large, and does not approach an integer either as $N \rightarrow \infty$. (A2)(b) guarantees that the interaction is not ‘conspiring’ to allow $|\sum_{i=\hat{m}+1}^n J_i N^i - h|$ to vanish as $N \rightarrow \infty$. Both assumptions are made to avoid certain degeneracies. These would not pose an essential problem, but would complicate our analysis unnecessarily.

Assumption (A3):

For all $1 \leq k \leq N^{\hat{m}}$ with N -ary decomposition $k = a_{\hat{m}-1} N^{\hat{m}-1} + \dots + a_0$:

$$(1.33) \quad \lim_{N \rightarrow \infty} \frac{\sum_{i=0}^{\hat{m}-1} J_{i+1} N^i \left[(N - a_i - 1) \left(\sum_{j=0}^i a_j N^j \right) + a_i \left(N^i - \sum_{j=0}^{i-1} a_j N^j \right) \right] + k \sum_{i=\hat{m}+1}^n J_i N^i}{[\hat{s}] (2\hat{s} - [\hat{s}] + 1) J_{\hat{m}+1} N^{2\hat{m}}} = 0.$$

This assumption has a somewhat unappealing form. Its purpose is to ensure that, in the limit as $N \rightarrow \infty$, the energy along optimal paths fluctuates by relatively small amounts over short distances. We will see that it is satisfied when $J_i = o(N^{-i+1})$ as $N \rightarrow \infty$.

Assumption (A4):

eq:A3decayrateofJ1

$$(1.34) \quad \frac{J_{i+1}}{J_i} = O\left(\frac{1}{N}\right) \quad \forall 1 \leq i \leq \hat{m}.$$

This assumption guarantees that the total interaction between a given spin and all the spins at a given hierarchical level remains bounded as $N \rightarrow \infty$.

Assumption (A5):

eq:AssumptionUniquemax

$$(1.35) \quad \text{No linear combination of } J_1, \dots, J_n \text{ is a multiple of } h.$$

This assumption again avoids certain degeneracies, and is valid for all but countably many choices of h and \vec{J} .

1.6. Main theorems. We are now ready to state our main results. The seven theorems and two corollaries given below identify the triple in (1.16), consisting of the communication height Γ^* , the set of critical configurations \mathcal{C}^* and the prefactor K^* . Formulas simplify as more constraints are placed on \vec{J} .

• **Communication height.** Recall the definition of Γ^* in (1.12).

Theorem 1.9. *Suppose that \vec{J} is non-increasing and that (A1) and (A3) hold. Then*

$$\text{eq:thrm-Gamma value} \quad (1.36) \quad \Gamma^* = [1 + o_N(1)] \frac{1}{4} (J_{\hat{m}+1})^{-1} \left(\sum_{i=\hat{m}+1}^n J_i N^i - h \right)^2, \quad N \rightarrow \infty.$$

Corollary 1.10. *Suppose that $J_i = \tilde{J}_i/N^i$ with $\tilde{J}_i = o(N)$ and that (A2)(b) holds. Then (A3) holds and*

$$(1.37) \quad \Gamma^* = [1 + o_N(1)] \frac{1}{4} (\tilde{J}_{\hat{m}+1})^{-1} \left(\sum_{i=\hat{m}+1}^n \tilde{J}_i - h \right)^2 N^{\hat{m}+1}, \quad N \rightarrow \infty.$$

Our next result gives a formula for Γ^* when $J_i = \tilde{J}/N^i$ for some $\tilde{J} > 0$. Let

$$\text{eq:indexI} \quad (1.38) \quad \mathbb{I} = \{(m, s) : 0 \leq m \leq n-1, 1 \leq s \leq N-1\} \cup \{(n-1, N)\} \subseteq \mathbb{N}^2,$$

and for $(m, s) \in \mathbb{I}$ define

$$\text{eq:hdagger} \quad (1.39) \quad h^{(m,s)} = \tilde{J} \left[\left(1 - \frac{1}{N}\right) (n-m) - (s-1) \frac{1}{N} \right] \in \left[0, \tilde{J} \left(1 - \frac{1}{N}\right) n\right].$$

Theorem 1.11. *Suppose that $J_i = \tilde{J}/N^i$ for some $\tilde{J} > 0$. Let $(m, s) \in \mathbb{I}$ be such that h satisfies*

$$(1.40) \quad h^{(m,s)} \leq h < h^{(m,s-1)}.$$

(1) *If N is odd, then*

$$\text{eq:thrm-Gamma-case3a} \quad (1.41) \quad \Gamma^* = \frac{\tilde{J}}{4N} \left[N^m \left(2s \left(N - \frac{s}{2} + s \bmod 2 \right) - N - s \bmod 2 \right) + N - 2s - (-1)^{s \bmod 2} \right] \\ + \frac{1}{2} \left[\tilde{J} \left(1 - \frac{1}{N} \right) (n-m-1) - h \right] (N^m (s - s \bmod 2) + 1).$$

(2) *If N is even and s is odd, then*

$$(1.42) \quad \Gamma^* = \Gamma_{m,s}^*$$

with

$$(1.43) \quad \Gamma_{m,s}^* = \frac{\tilde{J}}{2} N^{-m \bmod 2} (A_m - 1) + \tilde{J} \left[\frac{1}{2} B_m - N^{m \bmod 2} A_m \right] (N - s) \\ + \tilde{J} \left[\frac{N}{4} B_m - N^{m \bmod 2} A_m + N^{m-1} \left(\frac{s-1}{2} \right) \left(N - \frac{s-1}{2} \right) \right] \\ + \left[\left(\frac{s-1}{2} \right) N^m + \frac{N}{2} B_m - N^{1+m \bmod 2} A_m \right] \left[\tilde{J} \left(1 - \frac{1}{N} \right) (n-m-1) - h \right],$$

where $A_m = \left(\frac{N^{m-m \bmod 2} - 1}{N^2 - 1} \right)$ and $B_m = \left(\frac{N^m - 1}{N - 1} \right)$.

(3) *If N is even and s is even, then*

$$\text{eq:thrm-Gamma-case3c} \quad (1.44) \quad \Gamma^* = \Gamma_{m,s-1}^* + \left(h^{(m,s-1)} - h \right) \left[s N^m - \left(\frac{s-1}{2} \right) N^m - \left(\frac{N}{2} \right) B_m + N^{1+m \bmod 2} A_m \right].$$

Corollary 1.12. *Suppose that $J_i = \tilde{J}/N^i$ for some $\tilde{J} > 0$. Let $\alpha \in (0, 1)$ and $0 \leq m \leq n-1$ be such that $h = \tilde{J} (n-m-\alpha)$. Then*

$$\text{eq:thrm-Gamma-case4} \quad (1.45) \quad \Gamma^* = [1 + o_N(1)] \frac{\tilde{J}}{4} \alpha^2 N^{m+1}.$$

• **Critical configurations.** Recall the definition of \mathcal{C}^* in Definition 1.2. Recall from Section 1.4 that every integer $a \in \Lambda_N^n$ corresponds to a vertex v_a in such a way that $d(a, b) = d(v_a, v_b)$, and that $\gamma: \boxminus \rightarrow \boxplus$ is the reference path $\gamma = (\gamma_0, \dots, \gamma_{N^n})$, where γ_k is the configuration with $\gamma_k(v_a) = +1$ for $a \leq k$ and $\gamma_k(v_a) = -1$ for $a > k$. If \vec{J} is monotone, then γ is an optimal path as defined in (1.13).

Theorem 1.13. *Suppose that \vec{J} is strictly monotone. Then there exists a $1 \leq M \leq N^n$ such that \mathcal{C}^* is the set of isometric translations of γ_M . Furthermore, if (A1), (A2) and (A4) hold, then the N -ary decomposition $M = a_{n-1}N^{n-1} + \dots + a_0$ satisfies*

$$(1.46) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=0}^{n-1} |a_i - \eta_i| = 0,$$

where the coordinates $\eta_0, \dots, \eta_{n-1}$ are as follows: $\eta_i = 0$ for $\hat{m} < i \leq n-1$, $\eta_{\hat{m}} = \lceil \hat{s} \rceil$, and $\eta_{\hat{m}-1}, \dots, \eta_0$ are defined recursively in (3.28) and (3.32) below.

By isometric translation we mean any bijection $\phi: \Lambda_N^n \rightarrow \Lambda_N^n$ such that $d(v_a, v_b) = d(\phi(v_a), \phi(v_b))$, $1 \leq a, b \leq N^n$.

Theorem 1.14. *Suppose that \vec{J} is strictly monotone and that $J_i = \tilde{J}_i/N^i$ with $\tilde{J}_i = o(N)$. If (A1), (A2) and (A4) hold, then the coordinates $\eta_0, \dots, \eta_{n-1}$ in Theorem 1.13 are as follows:*

$$(1.47) \quad \eta_i = \begin{cases} 0, & \hat{m} < i \leq n-1, \\ \lceil \hat{s} \rceil, & i = \hat{m}, \\ \frac{N}{2}, & i = \hat{m}-1, \\ \frac{N}{2} \left[\sum_{j=1}^{i+1} \left(\frac{\tilde{J}_{\hat{m}-i+j}}{\tilde{J}_{\hat{m}-i}} \right) \left(1 - \frac{2\eta_{\hat{m}-i}}{N} \right) + \sum_{j=2}^{n-\hat{m}} \left(\frac{\tilde{J}_{\hat{m}+j}}{\tilde{J}_{\hat{m}-i}} \right) - \frac{h}{\tilde{J}_{\hat{m}-i}} + 1 \right], & 1 \leq i \leq \hat{m}-1. \end{cases}$$

Theorem 1.15. *Suppose that $J_i = \tilde{J}/N^i$ for some $\tilde{J} > 0$. Let $(m, s) \in \mathbb{I}$ be such that h satisfies*

$$(1.48) \quad h^{(m,s)} \leq h < h^{(m,s-1)}.$$

Then \mathcal{C}^* is the set of all isometric translations of the configuration γ_M , where

$$(1.49) \quad M = \begin{cases} \lceil \frac{s}{2} N^m \rceil, & N \text{ is odd and } s \text{ is odd,} \\ \lceil \frac{s-1}{2} N^m \rceil + 1, & N \text{ is odd and } s \text{ is even,} \\ \left(\frac{s-1}{2} N^m + \sum_{j=1}^{m-1} \left(\frac{N}{2} - (s+j+1) \bmod 2 \right) N^{m-j} + \frac{N}{2} \right), & N \text{ is even and } s \text{ is odd,} \\ \left(\frac{s-2}{2} N^m + \sum_{j=1}^{m-1} \left(\frac{N}{2} - (s+j+1) \bmod 2 \right) N^{m-j} + \frac{N}{2} \right), & N \text{ is even and } s \text{ is even.} \end{cases}$$

• **Prefactor.** We finally turn to the prefactor K^* defined in (1.17).

Theorem 1.16. *Suppose that \vec{J} is strictly monotone and that (A1)–(A5) hold. Then*

$$(1.50) \quad \frac{1}{K^*} = [1 + o_N(1)] \times \frac{\left[\sum_{i \in B_d} \eta_{i-1} N^{i-1} \right] \left[\sum_{i \in B_u} (N^i - \eta_{i-1} N^{i-1}) \right]}{\left[\sum_{i \in B_d} \eta_{i-1} N^{i-1} \right] + \left[\sum_{i \in B_u} (N^i - \eta_{i-1} N^{i-1}) \right]} \frac{N^{n-\hat{m}-1}}{N - \eta_0} \prod_{i=0}^{\hat{m}} \binom{N}{\eta_i} (N - \eta_i),$$

where $\eta_0, \dots, \eta_{n-1}$ are the coordinates defining the critical configurations in Theorem 1.13, and the integer sets B_d and B_u are defined in (3.39) below.

Theorem 1.17. *Suppose that $J_i = \tilde{J}/N^i$ for some $\tilde{J} > 0$ and that h satisfies*

$$(1.51) \quad h^{(m,s)} < h < h^{(m,s-1)}$$

for some $(m, s) \in \mathbb{I}$. If N is odd, $N \neq 2, 4$ and $m \geq 1$, then

$$(1.52) \quad \frac{1}{K^*} = a_0 N^{n-m-2} \prod_{i=0}^m \binom{N}{a_i} (N - a_i),$$

where $a_0 = \frac{N-1}{2} + 1$, $a_i = \frac{N-1}{2}$ for $i = 1, \dots, m-1$, and $a_m = \frac{s-1-(s+1)\bmod 2}{2}$.

1.7. Discussion. The theorems and corollaries in Section 1.6 provide a full description of the metastable behaviour of Glauber spin-flip dynamics on the hierarchical lattice, for any dimension N and any number of hierarchical levels n . The formulas are somewhat complicated for general \tilde{J} , but simplify considerably as more restrictions are imposed on \tilde{J} , such as $J_i = \tilde{J}/N^i$, $1 \leq i \leq n$ and $\tilde{J} > 0$, and in the hierarchical mean-field limit $N \rightarrow \infty$. The formulas even allow us to investigate the limit $n \rightarrow \infty$ towards the infinite hierarchical lattice.

The case of ‘standard’ interaction, defined by $J_i = \tilde{J}/N^i$ and treated in Section 4, is the easiest to interpret. The magnetic field h defines the integer pair (m, s) through the inequality

$$(1.53) \quad \tilde{J} \left[\left(1 - \frac{1}{N}\right) (n-m) - (s-1) \frac{1}{N} \right] \leq h < \tilde{J} \left[\left(1 - \frac{1}{N}\right) (n-m) - (s-2) \frac{1}{N} \right].$$

It turns out that the pair (m, s) captures the size of a critical configuration. Indeed, from Theorem 1.15 we see that if N is odd, then every critical configuration is of size $M = \lceil \frac{sN^m}{2} \rceil$ when s is odd and $M = \lceil \frac{(s-1)N^m}{2} \rceil$ when s is even, with similar results for N even. In particular, the set of critical configurations corresponds precisely to the set of all configurations of said size that are an isometric translation of γ_M .

Equations (1.41) and (1.44) in Theorem 1.11 are not particularly elegant, but in the hierarchical mean-field limit, and with $\alpha \in (0, 1)$ and $1 \leq m \leq n-1$ defined through the equation $h = \tilde{J}(n-m-\alpha)$, we find that

$$(1.54) \quad \lim_{N \rightarrow \infty} \frac{\Gamma^*}{N^{m+1}} = \frac{\tilde{J}\alpha^2}{4},$$

eq:Gamma limit

while for $\alpha = 0$ we have $\lim_{N \rightarrow \infty} \frac{\Gamma^*}{N^m} = \frac{1}{4}\tilde{J}$.

The prefactor K^* in Theorem 1.17 in the hierarchical mean-field limit scales like

$$(1.55) \quad \frac{1}{K^*} \sim \left(\frac{1-\alpha}{2} \right) 2^{m(N-\frac{1}{2})} N^n \binom{N}{\alpha N},$$

in which the dominant term is exponential in N .

Our results are part of a larger enterprise in which the goal is to understand metastability on large graphs. Jovanovski [4]^{pr} analysed the case of the *hypercube*, Dommers [2]^{pr} the case of the *random regular graph*, and Dommers, den Hollander, Jovanovski and Nardi [3]^{pr} the case of the *configuration model*. Each requires carrying out a detailed combinatorial analysis that is model-specific, even though the metastable behaviour expressed in Theorems 1.3–1.5 is universal. For lattices like the hypercube and the hierarchical lattice a full identification of the triple in (1.16) is possible, while for random graphs like the random regular graph and the configuration model so far only the communication height is well understood, while the set of critical configurations and the prefactor remain somewhat elusive.

2. MONOTONE PAIR POTENTIALS

In Section 2.1 we study the change in energy when all spins in two hierarchical blocks are switched (Lemma 2.1 below). In Section 2.2 we show that the reference path γ is an optimal paths for monotone pair potentials (Lemma 2.2 below). In Section 2.3 we give the proof of Theorem 1.8.

2.1. Energy landscape. Let $m \leq n-1$, let U be an $m+1$ -block in Λ_N^n , and let U_1 and U_2 be two disjoint m -blocks in U . Suppose that $U'_1 \subset U_1$ is a k -block in U_1 and $U'_2 \subset U_2$ is a k -block in U_2 , for some $k < m$. Let $\sigma \in \Omega$ be any configuration, and let σ' be the result of switching the values of σ at U'_1 and U'_2 . More precisely, let $\varphi: U'_1 \rightarrow U'_2$ be any isometric (with respect to d) bijection, and

set

$$(2.1) \quad \sigma'(v) = \begin{cases} \sigma(v), & v \notin U'_1 \cup U'_2, \\ \sigma(\varphi(v)), & v \in U'_1, \\ \sigma(\varphi^{-1}(v)), & v \in U'_2. \end{cases}$$

For $k+1 \leq i \leq m$, let $A_i = \{x \in U_1 \cap \bar{\sigma} : d(x, U'_1) = i\}$ (which is well defined because all $v \in U'_1$ are at the same distance from any $x \in U_1 \setminus U'_1$), $B_i = \{x \in U_1 \cap \sigma : d(x, U'_1) = i\}$, $C_i = \{x \in U_2 \cap \bar{\sigma} : d(x, U'_2) = i\}$ and $D_i = \{x \in U_2 \cap \sigma : d(x, U'_2) = i\}$.

Lemma 2.1. *For any $\sigma \in \Omega$,*

$$(2.2) \quad \mathcal{H}(\sigma') - \mathcal{H}(\sigma) = \sum_{i=k+1}^m 2(J_i - J_{m+1}) (|A_i| - |C_i|) (|U'_2 \cap \sigma| - |U'_1 \cap \sigma|).$$

Proof. Note that the number of interacting pairs (i.e., pairs (v, w) such that $\sigma(v) = -\sigma(w)$) in $U'_1 \times U'_2$ in σ is the same as in σ' . Hence

$$(2.3) \quad - \sum_{\substack{v \in U'_1 \\ w \in U'_2}} J_{d(v,w)} \sigma(v) \sigma(w) = - \sum_{\substack{v \in U'_1 \\ w \in U'_2}} J_{d(v,w)} \sigma'(v) \sigma'(w).$$

The same is true for interacting pairs in $(\overline{U'_1 \cup U'_2}) \times (\overline{U'_1 \cup U'_2})$, $U'_1 \times U'_1$, $U'_2 \times U'_2$, as well as $\bar{U} \times \Lambda_N^n$, where \bar{U} is the complement of U . Thus, we only need to consider interacting pairs coming from $U'_1 \times (U_1 \setminus U'_1)$, $U'_1 \times (U_2 \setminus U'_2)$, $U'_2 \times (U_2 \setminus U'_2)$ and $U'_2 \times (U_1 \setminus U'_1)$. The contribution to $\mathcal{H}(\sigma) - \mathcal{H}(\boxplus)$ of interacting pairs in $U'_1 \times (U_1 \setminus U'_1)$ is given by

$$(2.4) \quad - \sum_{\substack{v \in U'_1 \\ w \in U_1 \setminus U'_1}} J_{d(v,w)} \sigma(v) \sigma(w) = \sum_{i=k+1}^m J_i (|A_i| |U'_1 \cap \sigma| + |B_i| |U'_1 \cap \bar{\sigma}|).$$

Thus by moving the set $U'_1 \cap \sigma$ from U_1 to U_2 , this contribution is replaced by

$$(2.5) \quad - \sum_{\substack{v \in U'_2 \\ w \in U_1 \setminus U'_1}} J_{d(v,w)} \sigma'(v) \sigma'(w) = \sum_{i=k+1}^m J_{m+1} (|A_i| |U'_1 \cap \sigma| + |B_i| |U'_1 \cap \bar{\sigma}|).$$

Similarly, the contribution to $\mathcal{H}(\sigma) - \mathcal{H}(\boxplus)$ of interacting pairs in $U'_1 \times (U_2 \setminus U'_2)$ is given by

$$(2.6) \quad \sum_{i=k+1}^m J_{m+1} (|C_i| |U'_1 \cap \sigma| + |D_i| |U'_1 \cap \bar{\sigma}|),$$

which is subsequently replaced by

$$(2.7) \quad \sum_{i=k+1}^m J_i (|C_i| |U'_1 \cap \sigma| + |D_i| |U'_1 \cap \bar{\sigma}|).$$

Similar observations follow for interacting pairs in $U'_2 \times (U_2 \setminus U'_2)$ and $U'_2 \times (U_1 \setminus U'_1)$. Hence

$$(2.8) \quad \begin{aligned} \mathcal{H}(\sigma') - \mathcal{H}(\sigma) &= \sum_{i=k+1}^m (J_i - J_{m+1}) \\ &\times \left([|A_i| - |C_i|] (|U'_2 \cap \sigma| - |U'_1 \cap \sigma|) + [|B_i| - |D_i|] (|U'_2 \cap \bar{\sigma}| - |U'_1 \cap \bar{\sigma}|) \right). \end{aligned}$$

Noting that $|B_i| + |A_i| = (N-1)N^{i-1} = |D_i| + |C_i|$, we get

$$\begin{aligned}
\mathcal{H}(\sigma') - \mathcal{H}(\sigma) &= \sum_{i=k+1}^m (J_i - J_{m+1}) \\
&\times \left([|A_i| - |C_i|] (|U'_2 \cap \sigma| - |U'_1 \cap \sigma|) + [|C_i| - |A_i|] (|U'_2 \cap \bar{\sigma}| - |U'_1 \cap \bar{\sigma}|) \right) \\
(2.9) \quad &= \sum_{i=k+1}^m (J_i - J_{m+1}) \\
&\times \left([|A_i| - |C_i|] (|U'_2 \cap \sigma| - |U'_1 \cap \sigma|) + [|A_i| - |C_i|] (|U'_1 \cap \bar{\sigma}| - |U'_2 \cap \bar{\sigma}|) \right).
\end{aligned}$$

Finally, noting that $|U'_1 \cap \bar{\sigma}| = N^k - |U'_1 \cap \sigma|$ and $|U'_2 \cap \bar{\sigma}| = N^k - |U'_2 \cap \sigma|$, we complete the proof. \square

2.2. Optimal paths. Recall the definition of an optimal path from (1.13). We call a path $\gamma: \boxminus \rightarrow \boxplus$, denoted by $\{\gamma_i\}_{i=0}^M$ for some $M \geq N^n$, *uniformly optimal* when, for all $0 \leq i \leq M$,

eq:unif-optimal

$$(2.10) \quad \mathcal{H}(\gamma_i) = \min_{\substack{\sigma \in \Omega: \\ |\sigma| = |\gamma_i|}} \mathcal{H}(\sigma),$$

and *strictly optimal* when the minimum in the right-hand side of (1.13) is only attained by configurations that belong to some uniformly optimal path. We think of a path γ between two configurations in Ω both as a sequence of configurations denoted by $\{\gamma_i\}_{i=1}^M$ and as a sequence of vertices denoted by $\{\gamma(i)\}_{i=1}^M$, where $\gamma(i)$ is the single vertex in the symmetric difference $\gamma_{i-1} \Delta \gamma_i$.

Order the vertices $\{v_i\}_{i=1}^{N^n}$ in Λ_N^n in a natural order so that, for all $1 \leq k \leq n-1$ and for all $0 \leq j \leq N^n/N^k$, $\{v_{jN^k+1}, \dots, v_{(j+1)N^k}\}$ belong to the same k -block. Let $\gamma^{\text{MD}}: \boxminus \rightarrow \boxplus$ be the path defined by $\gamma^{\text{MD}}(i) = v_i$ for $1 \leq i \leq N^n$. Let $\gamma^{\text{MI}}: \boxminus \rightarrow \boxplus$ be defined by $\gamma^{\text{MI}}(k) = v_{\theta(k)}$ and

$$(2.11) \quad \theta(k) = 1 + \sum_{i=0}^{n-1} N^{n-1-i} \left(\left\lfloor \frac{k-1}{N^i} \right\rfloor \bmod N \right).$$

Thus, the vertex $\gamma^{\text{MI}}(k)$ belongs to the $((k-1) \bmod N)^{\text{th}}$ $(n-1)$ -block, and within that block it belongs to the $(\lfloor \frac{k-1}{N^2} \rfloor \bmod N)^{\text{th}}$ $(n-2)$ -block, etc. We can now use Lemma 2.1 to draw the following conclusions.

- Lemma 2.2.** (1) If \vec{J} is non-increasing, then γ^{MD} is a uniformly optimal path.
(2) If \vec{J} is non-decreasing, then γ^{MI} is a uniformly optimal path.
(3) If \vec{J} is strictly decreasing or strictly increasing, then γ^{MD} or γ^{MI} is strictly optimal.

Proof. We treat the non-increasing case and the non-decreasing case separately.

Non-increasing case: Let $\sigma \in \Omega$ be given. We will construct a sequence of configurations $\{\psi_i\}_{i=1}^n$, all of volume $|\sigma|$ and with $\psi_n = \gamma^{\text{MD}}_{|\sigma|}$, such that $\mathcal{H}(\sigma) \geq \mathcal{H}(\psi_1) \geq \dots \geq \mathcal{H}(\psi_n)$, and the inequalities being strict whenever \vec{J} is strictly decreasing. This will prove the claim for the non-increasing case.

For $1 \leq k \leq n$, define ψ_k to be the (unique) configuration that satisfies the following two conditions:

1. For every k -block $U \subset \Lambda_N^n$, $|U \cap \sigma| = |U \cap \psi_k|$.
2. For all $i < j$ with v_i and v_j belonging in the same k -block, $v_j \in \psi_k$ implies $v_i \in \psi_k$.

In particular, note that ψ_1 is obtained from σ by “shifting” the $+1$ values of σ found inside every 1-block as far left as possible (i.e., with the lowest possible index) within the same 1-block. It is obvious that $\mathcal{H}(\psi_1) = \mathcal{H}(\sigma)$. It is also clear from this recursive definition that $\psi_n = \gamma^{\text{MD}}_{|\sigma|}$.

Starting with ψ_k , we will show how to obtain ψ_{k+1} by a series of transformations that are non-increasing in \mathcal{H} . Let U be the first $k+1$ block of Λ_n^N , and let U_1, \dots, U_N be its k -blocks,

arranged so that $|U_i \cap \sigma| \geq |U_{i+1} \cap \sigma|$. Note that this may be achieved by re-arranging (or re-labeling) the blocks U_1, \dots, U_N , and any such re-arranging is an \mathcal{H} -preserving operation. Let $a = \min \{i: |U_i \cap \sigma| < N^k\}$ and $b = \max \{i: |U_i \cap \sigma| > 1\}$. Note that if $a = b$, then $U \cap \sigma$ is already in the correct form, satisfying the definition of ψ_{k+1} . Thus, we may assume that $a \neq b$. Find a maximal block $\tilde{U}_b \subsetneq U_b$ with $|\tilde{U}_b \cap \sigma| > 0$ such that, for some block of equal size $\tilde{U}_a \subsetneq U_a$, $|\tilde{U}_b \cap \sigma| > |\tilde{U}_a \cap \sigma|$. To do this, take the first $k-1$ -block U'_b in U_b and the last $k-1$ -block U'_a in U_a that satisfies $|U'_a \cap \sigma| > 0$, and check whether $|U'_b \cap \sigma| > |U'_a \cap \sigma|$. If not, then proceed by taking the first $k-2$ -block in U_b , etc. By the definition of a and b , this constructive search for \tilde{U}_a and \tilde{U}_b always yields two such blocks. Once these are found, perform the switching operation in Lemma 2.1 on the blocks \tilde{U}_a and \tilde{U}_b , and denote the resulting configuration by ψ'_k (see Fig. 6). Then, by Lemma 2.1, with s denoting the size of the blocks \tilde{U}_a and \tilde{U}_b ,

$$(2.12) \quad \mathcal{H}(\psi'_k) - \mathcal{H}(\psi_k) = \sum_{i=s+1}^k 2(J_i - J_{k+1}) [|A_i| - |C_i|] (|\tilde{U}_b \cap \sigma| - |\tilde{U}_a \cap \sigma|),$$

where we recall that $A_i = \{x \in U_a \cap \bar{\sigma}: d(x, \tilde{U}_a) = i\}$ and $C_i = \{x \in U_b \cap \bar{\sigma}: d(x, \tilde{U}_b) = i\}$. By definition, we have $|\tilde{U}_b \cap \sigma| - |\tilde{U}_a \cap \sigma| > 0$, and from the monotonicity we get that $J_i - J_{m+1} \geq 0$. Lastly, by the fact that $|U_a \cap \sigma| \geq |U_b \cap \sigma|$ and the construction of ψ_k , as well as the definition of \tilde{U}_b and \tilde{U}_a , it also follows that $|A_i| - |C_i| \leq 0$ for all $s+1 \leq i \leq k$. Therefore $\mathcal{H}(\psi'_k) - \mathcal{H}(\psi_k) \leq 0$. Repeating this construction until $\min \{i: |U_i \cap \sigma| < N^k\} = \max \{i: |U_i \cap \sigma| > 1\}$ (which happens in a finite number of moves), and repeating the same construction for all other $k+1$ -blocks, we get the configuration ψ_{k+1} , and hence $\mathcal{H}(\psi_{k+1}) - \mathcal{H}(\psi_k) \leq 0$.




FIGURE 6. The transformation $\psi_k \rightarrow \psi'_k$. The blocks \tilde{U}_a and \tilde{U}_b are drawn with a dashed outline. Solid black circles represent elements of ψ_k (i.e., vertices on which the configuration ψ_k takes the value $+1$), while blank circles are elements of $\bar{\psi}_k$.

Non-decreasing case: Given a configuration σ , we again apply a series of transformations involving switching and re-arranging of blocks in σ (all of which are non-increasing in \mathcal{H}) and ending with the configuration $\gamma_{|\sigma|}^{\text{MI}}$. Firstly, through a series of re-arrangements, we may assume that σ is *left-aligned*: for any $0 \leq k \leq n-1$ and any k -blocks U_i and U_{i+1} contained in the same $(k+1)$ -block (a lower index on a block implies that it contains vertices that also have a lower index), we have $|U_i \cap \sigma| \geq |U_{i+1} \cap \sigma|$. It is clear that these re-arrangements are \mathcal{H} -invariant.

Start with $k = n-1$ and check whether $|U_1 \cap \sigma| \geq |U_N \cap \sigma| + 2$. If so, then switch the value at $v_1 \in U_1$ (equal to $+1$) with the value at $v_N \in U_N$ (equal to -1). Denote the result of this switch by σ' . From Lemma 2.1 we have

$$(2.13) \quad \mathcal{H}(\sigma') - \mathcal{H}(\sigma) = \sum_{i=1}^{n-1} 2(J_i - J_n) [|A_i| - |C_i|] (0 - 1).$$

Since σ is left-aligned, we know that $|A_{n-1}| \leq |C_{n-1}|$. Inductively it follows that $|A_i| \leq |C_i|$ for all $1 \leq i \leq n-1$. Since, by the monotonicity, we also have $J_i - J_n \leq 0$ for all $1 \leq i \leq n-1$, it follows that $\mathcal{H}(\sigma') - \mathcal{H}(\sigma) \leq 0$.

Next re-arrange σ' to make it left-aligned (at no cost in \mathcal{H}), and repeat this construction until $|U_N \cap \sigma| \leq |U_1 \cap \sigma| \leq |U_N \cap \sigma| + 1$. Note that this takes a finite number of steps. Once this is

accomplished, resume by recursively repeating the construction for $k = n - 2$, within each $n - 1$ -block, etc. This terminates with $\gamma_{|\sigma|}^{\text{MI}}$. \square

2.3. Proof of Theorem 1.8. The proof is analogous to that given in [1, Section 17.3.1]^{BdH15}, and relies on the existence of a uniformly optimal path.

Proof. Let $\sigma \in \Omega \setminus \{\boxminus, \boxplus\}$. Find two vertices $v_i, v_j \in \Lambda_N^n$ such that $v_i \in \sigma$ and $v_j \notin \sigma$. By translation invariance, we can construct a uniformly optimal reference path γ that is a translation (via some d -preserving bijection of Λ_N^n) of the path γ^{MD} in the non-increasing case and γ^{MI} in the non-decreasing case, and that satisfies $\gamma(1) = v_j$ and $\gamma(2) = v_i$. Note that in both cases

$$\begin{aligned} \text{eq:intersectedpath} \quad (2.14) \quad & \sigma \cap \gamma_1 = \boxminus, \\ & 1 \leq |\sigma \cap \gamma_k| < k \quad \forall k \geq 2. \end{aligned}$$

Furthermore,

$$\text{eq:refpathcompare} \quad (2.15) \quad \mathcal{H}(\sigma \cup \gamma_1) - \mathcal{H}(\sigma) = \sum_{\substack{w \neq v_j \\ w \notin \sigma}} J_{d(w, v_j)} - \sum_{\substack{w \neq v_j \\ w \in \sigma}} J_{d(w, v_j)} - h < \sum_{w \neq v_j} J_{d(w, v_j)} - h = \mathcal{H}(\gamma_1) - \mathcal{H}(\boxminus)$$

where we use the fact that $J_i > 0$, $1 \leq i \leq n$. Similarly, if we let $k' = \min \{k \in \mathbb{N} : \mathcal{H}(\gamma_k) \leq \mathcal{H}(\boxminus)\}$, then by (A1) it follows that $k' \geq 2$, and so for $2 \leq k \leq k'$,

$$\begin{aligned} \text{eq:refpath} \quad (2.16) \quad & \mathcal{H}(\sigma \cup \gamma_k) - \mathcal{H}(\sigma) = \sum_{w \in \gamma_k \setminus \sigma} \sum_{v \notin \sigma \cup \gamma_k} J_{d(w, v)} - \sum_{w \in \gamma_k \setminus \sigma} \sum_{v \in \sigma} J_{d(w, v)} - h |\gamma_k \setminus \sigma| \\ & \leq \sum_{w \in \gamma_k \setminus \sigma} \sum_{v \notin \gamma_k} J_{d(w, v)} - \sum_{w \in \gamma_k \setminus \sigma} \sum_{v \in \sigma \cap \gamma_k} J_{d(w, v)} - h |\gamma_k \setminus \sigma| \\ & = \mathcal{H}(\gamma_k) - \mathcal{H}(\gamma_k \cap \sigma) \leq \mathcal{H}(\gamma_k) - \mathcal{H}(\gamma_{|\gamma_k \cap \sigma|}) < \mathcal{H}(\gamma_k) - \mathcal{H}(\boxminus), \end{aligned}$$

where the last inequality follows from the fact that $|\gamma_k \cap \sigma| < k$ (by (2.14)) because γ is uniformly optimal. Taking $k = k'$, we get from (2.16) that $\mathcal{H}(\sigma \cup \gamma_{k'}) < \mathcal{H}(\sigma)$, and hence that the stability level V_σ of σ defined in 1.8 satisfies

$$(2.17) \quad V_\sigma < \max_{1 \leq k \leq k'} \{0, (\mathcal{H}(\gamma_k) - \mathcal{H}(\boxminus))\} \leq \Gamma^*.$$

This settles the claim because $V_{\boxminus} = \Gamma^*$. \square

Remark 2.3. Note that if (A1) is not satisfied, or in other words if

$$\text{eq:nometastability} \quad (2.18) \quad \left(1 - \frac{1}{N}\right) \sum_{i=1}^n J_i N^i \leq h,$$

then it follows from the inequality in 2.15 (note that without (A1) this is not a strict inequality) that

$$(2.19) \quad \mathcal{H}(\sigma \cup \gamma_1) - \mathcal{H}(\sigma) \leq \mathcal{H}(\gamma_1) - \mathcal{H}(\boxminus) \leq 0,$$

and hence σ is not a local minimum of \mathcal{H} . Since σ is arbitrary, it follows that \mathcal{H} has no local minima. This again illustrates why assumption (A1) is needed.

3. NON-INCREASING PAIR POTENTIAL

In Section 3.1 we prove a concavity property for the energy profile along the reference path inside hierarchical blocks (Lemma 3.1 below). In Section 3.2 we show that the fluctuations of the energy profile inside a hierarchical block are relatively small (Lemma 3.2 below) and use this to prove Theorem 1.9 in the hierarchical mean-field limit (Corollary 3.3 and Remark 3.4 below). In Section 3.3 we identify the critical configurations and check that the conditions in Lemma 1.6 are satisfied (Lemmas 3.5–3.6 below). We use these results in Section 3.4 to prove Theorem 1.16 and in Section 3.5 to prove Theorems 1.13–1.14.

3.1. Concavity along the reference path. From now on we will only consider the case where \vec{J} is non-increasing. We will drop the superscript MD and denote the uniformly optimal path γ^{MD} defined in Section 2 by γ . We observe that

$$(3.1) \quad \mathcal{H}(\gamma_k) - \mathcal{H}(\Xi) = \sum_{i=1}^k \sum_{j=k+1}^{N^n} J_{d(v_i, v_j)} - hk, \quad 1 \leq k \leq N^n,$$

and it is not difficult to show that (3.1) can be written as

$$(3.2) \quad \mathcal{H}(\gamma_k) - \mathcal{H}(\Xi) = \sum_{i=1}^n J_i N^{i-1} \left(k \bmod N^i \left(N - \left\lfloor \frac{k}{N^{i-1}} \right\rfloor \bmod N - 1 \right) \right. \\ \left. + (N^{i-1} - k \bmod N^{i-1}) \left\lfloor \frac{k}{N^{i-1}} \right\rfloor \bmod N \right) - hk.$$

Hence the communication height between Ξ and Θ is given by

$$(3.3) \quad \Gamma^* = \max_{1 \leq k \leq N^n} \left\{ \sum_{i=1}^n J_i N^{i-1} \left(k \bmod N^i \left(N - \left\lfloor \frac{k}{N^{i-1}} \right\rfloor \bmod N - 1 \right) \right. \right. \\ \left. \left. + (N^{i-1} - k \bmod N^{i-1}) \left\lfloor \frac{k}{N^{i-1}} \right\rfloor \bmod N \right) - hk \right\}.$$

However, it is not clear from (3.3) how Γ^* and the energy values along the path γ depend on \vec{J} . We will therefore derive Γ^* in a different way, obtaining a more insightful expression.

Note that if $j < k$, then

$$(3.4) \quad \mathcal{H}(\gamma_k) - \mathcal{H}(\gamma_j) = \sum_{i=j+1}^k \left(\sum_{s=k+1}^{N^n} J_{d(v_i, v_s)} - \sum_{s=1}^j J_{d(v_i, v_s)} \right) - h(k-j).$$

In particular, we observe that, for any $0 \leq a \leq n-1$,

$$(3.5) \quad \mathcal{H}(\gamma_{N^a}) - \mathcal{H}(\gamma_0) = \mathcal{H}(\gamma_{N^a}) - \mathcal{H}(\Xi) = (N-1)N^a \sum_{i=a}^{n-1} N^i J_{i+1} - hN^a.$$

We are interested in the global maxima of the energy profile. In order to locate where these occur, we analyse the geometric properties of the sequence $\{\mathcal{H}(\gamma_i)\}_{i=0}^{N^n}$. The following result describes concave subsequences that appear in $\{\mathcal{H}(\gamma_i)\}_{i=0}^{N^n}$ (see Fig. 7) and that will be used repeatedly in Section 4 to locate the global maxima of the energy landscape.

Lemma 3.1. *Suppose that $k = j + N^a$ and $l = k + N^a$ for some $a \geq 0$ and $j \geq 0$. Suppose that the three vertices v_j , v_k and v_l all lie in the same $(a+1)$ -block. Then*

$$(3.6) \quad (\mathcal{H}(\gamma_k) - \mathcal{H}(\gamma_j)) - (\mathcal{H}(\gamma_l) - \mathcal{H}(\gamma_k)) = 2J_{a+1}N^{2a}.$$

Proof. Note that, for any $1 \leq s \leq N^a$, $b \geq 1$, $b \neq a+1$,

$$(3.7) \quad |\{t > j + N^a : d(v_{j+s}, v_t) = b\}| = |\{t > k + N^a : d(v_{k+s}, v_t) = b\}|,$$

while

$$(3.8) \quad |\{t > j + N^a : d(v_{j+s}, v_t) = a+1\}| = |\{t > k + N^a : d(v_{k+s}, v_t) = a+1\}| + N^a.$$

Similarly, for $b \geq 1$, $b \neq a+1$,

$$(3.9) \quad |\{t \leq j : d(v_{j+s}, v_t) = b\}| = |\{t \leq k : d(v_{k+s}, v_t) = b\}|,$$

while

$$(3.10) \quad |\{t \leq j : d(v_{j+s}, v_t) = a+1\}| + N^a = |\{t \leq k : d(v_{k+s}, v_t) = a+1\}|.$$

Hence, by rewriting the sum in (3.4), we get

$$\begin{aligned}
& (\mathcal{H}(\gamma_k^{\text{MD}}) - \mathcal{H}(\gamma_j^{\text{MD}})) - (\mathcal{H}(\gamma_i^{\text{MD}}) - \mathcal{H}(\gamma_k^{\text{MD}})) \\
&= \left(\sum_{s=1}^{N^a} \sum_{b=1}^n J_b |\{t > j + N^a : d(v_{j+s}, v_t) = b\}| - \sum_{s=1}^{N^a} \sum_{b=1}^n J_b |\{t \leq j : d(v_{j+s}, v_t) = b\}| \right) \\
& \quad - \left(\sum_{s=1}^{N^a} \sum_{b=1}^n J_b |\{t > k + N^a : d(v_{k+s}, v_t) = b\}| - \sum_{s=1}^{N^a} \sum_{b=1}^n J_b |\{t \leq k : d(v_{k+s}, v_t) = b\}| \right) \\
&= 2J_{a+1}N^{2a}.
\end{aligned}
\tag{3.11}$$

This shows that the energy profile along the path γ is made up of periodic segments that are *concave* (see Definition 4.1 below). \square




FIGURE 7. The solid circles represent a periodic subsequence of $\{\mathcal{H}(\gamma_i)\}_{i=0}^{N^n}$ of period N^{n-1} , while the hollow circles represent points of period N^{n-2} that are contained within the same $(n-1)$ -block.

3.2. Hierarchical mean-field limit. The hierarchical mean-field limit corresponds to letting the hierarchical dimension N tend to infinity while keeping the hierarchical height n fixed. We will show that, under certain assumptions on the rate of decay of the sequence $\{J_i\}_{i=1}^n$, in the hierarchical mean-field limit the sequence $\{\mathcal{H}(\gamma_i)\}_{i=0}^{N^n}$ attains its global maximum at a location that is close to a multiple (by some factor in $\{1, \dots, N\}$) of the largest block size where the corresponding configuration has energy larger than $\mathcal{H}(\boxminus)$. We define this explicitly as follows.

Recall from (1.31) that

$$\begin{aligned}
\hat{m} &= \max \left\{ 0 \leq m \leq n-1 : \left(1 - \frac{1}{N}\right) \sum_{i=m+1}^n J_i N^i > h \right\} \\
(3.12) \quad &= \max \{ 0 \leq m \leq n-1 : \mathcal{H}(\gamma_{N^m}) \geq \mathcal{H}(\boxminus) \},
\end{aligned}$$

where the second line follows from (3.5).

From Lemma 3.1 it follows that, for all $M > \hat{m}$ and all $1 \leq s \leq N-1$, $\mathcal{H}(\gamma_{sN^M}) < \mathcal{H}(\Theta)$. Note also that, by Lemma 3.1 and equation (3.5), we define

$$\begin{aligned}
\alpha_{\hat{m},s} &= \mathcal{H}(\gamma_{sN^{\hat{m}}}) - \mathcal{H}(\Theta) \\
&= \sum_{i=0}^{s-2} (\mathcal{H}(\gamma_{(s-i)N^{\hat{m}}}) - \mathcal{H}(\gamma_{(s-i-1)N^{\hat{m}}})) + \mathcal{H}(\gamma_{N^{\hat{m}}}) - \mathcal{H}(\gamma_0) \\
&= s \left[(\mathcal{H}(\gamma_{N^{\hat{m}}}) - \mathcal{H}(\gamma_0)) - (s-1) J_{\hat{m}+1} N^{2\hat{m}} \right] \\
(3.13) \quad &= sN^{\hat{m}} \left[\left(1 - \frac{1}{N}\right) \sum_{k=\hat{m}}^{n-1} J_{k+1} N^{k+1} - h - (s-1) J_{\hat{m}+1} N^{\hat{m}} \right].
\end{aligned}$$

Increments of values given by (3.13) are equal to

$$(3.14) \quad \alpha_{\hat{m},s+1} - \alpha_{\hat{m},s} = N^{\hat{m}} \left[\left(1 - \frac{1}{N}\right) \sum_{k=\hat{m}}^{n-1} J_{k+1} N^{k+1} - h - 2s J_{\hat{m}+1} N^{\hat{m}} \right].$$

By the concavity implied by Lemma 3.1, we have that $\alpha_{\hat{m},s+1} - \alpha_{\hat{m},s} \leq 0$ if and only if $s \geq \hat{s}$, where \hat{s} is defined in (1.32). Under Assumption (A1)(a) it is easy to see that the sequence $\{\mathcal{H}(\gamma_{sN^{n-1}}) - \mathcal{H}(\Theta)\}_{s=0}^N$ attains a unique maximum at $1 \leq [\hat{s}] < N$, with value

$$(3.15) \quad \mathcal{H}(\gamma_{[\hat{s}]N^{\hat{m}}}) - \mathcal{H}(\Theta) = [\hat{s}] (2\hat{s} - [\hat{s}] + 1) J_{\hat{m}+1} N^{2\hat{m}}.$$

Furthermore, we claim that for any $N < t \leq N^{n-\hat{m}}$, $\mathcal{H}(\gamma_{tN^{\hat{m}}}) < \mathcal{H}(\gamma_{[\hat{s}]N^{\hat{m}}})$. Indeed, define $\bar{d} = d(v_{[\hat{s}]N^{\hat{m}}}, v_{tN^{\hat{m}}}) > \hat{m}$, and note that $tN^{\hat{m}} = \eta N^{\bar{d}} + sN^{\hat{m}}$ for some $0 \leq \eta, s < N$. Hence

$$\begin{aligned}
\mathcal{H}(\gamma_{tN^{\hat{m}}}) - \mathcal{H}(\Theta) &= \mathcal{H}(\gamma_{\eta N^{\bar{d}}}) - \mathcal{H}(\Theta) + \mathcal{H}(\gamma_{tN^{\hat{m}}}) - \mathcal{H}(\gamma_{\eta N^{\bar{d}}}) \\
&\leq \mathcal{H}(\gamma_{tN^{\hat{m}}}) - \mathcal{H}(\gamma_{\eta N^{\bar{d}}}) \\
(3.16) \quad &= sN^{\hat{m}} \left[\left(1 - \frac{1}{N}\right) \sum_{k=\hat{m}+1}^n J_k N^k - h - (s-1) J_{\hat{m}+1} N^{\hat{m}} - \eta J_{\bar{d}+1} N^{\bar{d}} \right] \\
&< \mathcal{H}(\gamma_{sN^{\hat{m}}}) - \mathcal{H}(\Theta) \leq \mathcal{H}(\gamma_{[\hat{s}]N^{\hat{m}}}) - \mathcal{H}(\Theta),
\end{aligned}$$

where the first inequality follows from the definition of \hat{m} and the fact that $\bar{d} > \hat{m}$.

We next show that fluctuations in energy $|\mathcal{H}(\gamma_i) - \mathcal{H}(\gamma_j)|$ for $|i-j| \leq N^{\hat{m}}$ are relatively small compared to $\mathcal{H}(\gamma_{[\hat{s}]}) - \mathcal{H}(\Theta)$.

Lemma 3.2. *Let $k = \sum_{i=0}^s a_i N^i$ with $0 \leq a_i \leq N-1$, and let $M = \sum_{i=t}^{n-1} a_i N^i$ with $0 \leq b_i \leq N-1$ and $n-1 \geq t > s$. Then*

$$(3.17) \quad \mathcal{H}(\gamma_{M+k}) - \mathcal{H}(\gamma_M) \leq \mathcal{H}(\gamma_k) - \mathcal{H}(\Theta)$$

and

$$(3.18) \quad |\mathcal{H}(\gamma_{M+k}) - \mathcal{H}(\gamma_M)| \leq |\mathcal{H}(\gamma_k) - \mathcal{H}(\Theta)| + hk.$$

Proof. Note that, during the move from γ_M to γ_{M+k} , the total change in energy due to interacting pairs at distance i is given by $(1 - \frac{1}{N}) k \sum_{i=s+2}^t J_i N^i$ for $s+2 \leq i \leq t$, while for $i \geq t+1$ it is given by $k \sum_{i=t}^{n-1} J_{i+1} N^i (N - 2b_i - 1)$. Now, for $1 \leq i \leq s+1$, this change is equal to

$$(3.19) \quad J_1 N^0 a_0 (N - a_0) + \sum_{i=1}^s J_{i+1} N^i \left((N - a_i - 1) \left(\sum_{j=0}^i a_j N^j \right) + a_i \left(N^i - \sum_{j=0}^{i-1} a_j N^j \right) \right),$$

which is also the same during the move from γ_{\boxminus} to γ_k . Thus, we get

$$\begin{aligned}
& \mathcal{H}(\gamma_{M+k}) - \mathcal{H}(\gamma_M) \\
&= \sum_{i=0}^s J_{i+1} N^i \left((N - a_i - 1) \left(\sum_{j=0}^i a_j N^j \right) + a_i \left(N^i - \sum_{j=0}^{i-1} a_j N^j \right) \right) \\
&\quad + \left(1 - \frac{1}{N} \right) k \sum_{i=s+2}^t J_i N^i + k \sum_{i=t}^{n-1} J_{i+1} N^i (N - 2b_i - 1) - hk \\
&\leq \sum_{i=0}^s J_{i+1} N^i \left((N - a_i - 1) \left(\sum_{j=0}^i a_j N^j \right) + a_i \left(N^i - \sum_{j=0}^{i-1} a_j N^j \right) \right) \\
&\quad + \left(1 - \frac{1}{N} \right) k \sum_{i=s+2}^n J_i N^i - hk \\
&= \mathcal{H}(\gamma_k) - \mathcal{H}(\gamma_{\boxminus}).
\end{aligned}
\tag{3.20}$$

Note, furthermore, that the right-hand side of the first line of (3.20) is non-negative, as is the first sum in the second line and both sums in the third line. Making use of the triangle inequality, we get the second claim of the lemma. \square

We will assume for now that $\hat{m} \geq 1$ and consider the case $\hat{m} = 0$ in Remark 3.4. It follows from Lemma 3.2 and Assumption (A3) that, for any $0 \leq k < N^{\hat{m}}$ and $\ell \geq 1$,

$$\frac{|\mathcal{H}(\gamma_{k+\ell N^{\hat{m}}}) - \mathcal{H}(\gamma_{\ell N^{\hat{m}}})|}{|\mathcal{H}(\gamma_{\lceil \hat{s} \rceil N^{\hat{m}}}) - \mathcal{H}(\boxminus)|} \leq \frac{|\mathcal{H}(\gamma_k) - \mathcal{H}(\gamma_{\boxminus})| + hk}{|\mathcal{H}(\gamma_{\lceil \hat{s} \rceil N^{\hat{m}}}) - \mathcal{H}(\boxminus)|} \rightarrow 0 \quad \text{as } N \rightarrow \infty,
\tag{3.21}$$

since from (3.20) we see that the numerator in the right-hand side of (3.21) equals the numerator in the condition of Assumption (A3), and from (3.13) the same follows for the denominator. Thus, using (3.13) we conclude the following.

Corollary 3.3 (Proof of Theorem 1.9). *Suppose that Assumption (A2) holds. Then*

$$\begin{aligned}
\Gamma^* &= [1 + o_N(1)] (\mathcal{H}(\gamma_{\lceil \hat{s} \rceil N^{\hat{m}}}) - \mathcal{H}(\boxminus)) \\
&= [1 + o_N(1)] \lceil \hat{s} \rceil (2\hat{s} - \lceil \hat{s} \rceil + 1) J_{\hat{m}+1} N^{2\hat{m}} \\
&= [1 + o_N(1)] \hat{s}^2 J_{\hat{m}+1} N^{2\hat{m}}.
\end{aligned}
\tag{3.22}$$

Remark 3.4. The special case $\hat{m} = 0$ can be considered separately. By Lemma 3.2 it follows, for any $0 \leq t \leq N^n$ and with

$$\hat{s} = (2J_1)^{-1} \left[\left(1 - \frac{1}{N} \right) \sum_{i=0}^{n-1} J_{i+1} N^{i+1} - h \right],
\tag{3.23}$$

that

$$\mathcal{H}(\gamma_t) - \mathcal{H}(\boxminus) \leq \mathcal{H}(\gamma_{\lceil \hat{s} \rceil}) - \mathcal{H}(\boxminus)
\tag{3.24}$$

and hence $\Gamma^* = \mathcal{H}(\gamma_{\lceil \hat{s} \rceil}) - \mathcal{H}(\boxminus) = \lceil \hat{s} \rceil (2\hat{s} - \lceil \hat{s} \rceil + 1) J_1$.

3.3. Critical configurations. It is clear from (1.17) that the prefactor K^* is closely related to the set of critical configurations \mathcal{C}^* , in particular, the cardinality of this set. The symmetry of Λ_N^n implies that the image of any critical configuration under an isometric translation is also a critical configuration. Thus, we have to count the number of isometries that result in distinct elements of \mathcal{C}^* , which is a problem related to the N -ary decomposition of the size of a critical configuration. To do so, we first establish a result that determines the N -ary decomposition of any global maximum subject to Assumption (A3).

The following lemma gives us the asymptotic value of the terms in the N -ary decomposition of the size of a critical configuration.

Lemma 3.5. *Suppose that (A1)–(A4) holds, and that the path γ attains a global maximum at γ_M . Let*

$$(3.25) \quad M = a_{n-1}N^{n-1} + \dots + k_1N + a_0$$

be the N -ary decomposition of the integer M . Then

$$(3.26) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=0}^{n-1} |a_i - \eta_i| = 0,$$

where $\eta_i = 0$ for $\hat{m} < i \leq n-1$, $\eta_{\hat{m}} = \lceil \hat{s} \rceil$, and $\eta_{\hat{m}-1}, \dots, \eta_0$ are defined in (3.28) and (3.32) below.

Proof. From the definition of \hat{m} in (1.31) and the argument leading up to (3.22), it is clear that $\lim_{N \rightarrow \infty} |a_i| = 0$ for $i > \hat{m}$. Let $\eta_{\hat{m}} = \lceil \hat{s} \rceil$. Then, for any $0 \leq \sigma < N$,

$$\begin{aligned} & \mathcal{H}(\gamma_{\eta_{\hat{m}}N^{\hat{m}} + \sigma N^{\hat{m}-1}}) - \mathcal{H}(\gamma_{\eta_{\hat{m}}N^{\hat{m}}}) \\ &= J_{\hat{m}}N^{2\hat{m}-2}\sigma(N-\sigma) + J_{\hat{m}+1}N^{2\hat{m}-1}\sigma(N-\eta_{\hat{m}}-1) \\ & \quad + \left(1 - \frac{1}{N}\right) \sum_{i=\hat{m}+2}^n \sigma J_i N^{\hat{m}-1+i} - J_{\hat{m}+1}\eta_{\hat{m}}\sigma N^{2\hat{m}-1} - h\sigma N^{\hat{m}-1} \\ (3.27) \quad &= J_{\hat{m}}N^{2\hat{m}-2}\sigma(N-\sigma) + J_{\hat{m}+1}N^{2\hat{m}-1}\sigma(N-2\eta_{\hat{m}}-1) \\ & \quad + \left(1 - \frac{1}{N}\right) \sum_{i=\hat{m}+2}^n \sigma J_i N^{\hat{m}-1+i} - h\sigma N^{\hat{m}-1}. \end{aligned}$$

By the concavity in Lemma 3.1, $\mathcal{H}(\gamma_{\eta_{\hat{m}}N^{\hat{m}} + (\sigma+1)N^{\hat{m}-1}}) - \mathcal{H}(\gamma_{\eta_{\hat{m}}N^{\hat{m}} + \sigma N^{\hat{m}-1}}) \leq 0$ if and only if

$$(3.28) \quad 0 \vee \left[\frac{1}{2} \left(\left(\frac{J_{\hat{m}+1}}{J_{\hat{m}}} \right) N(N-2\eta_{\hat{m}}-1) + \left(1 - \frac{1}{N}\right) \sum_{i=2}^{n-\hat{m}} \left(\frac{J_{\hat{m}+i}}{J_{\hat{m}}} \right) N^{i+1} - \frac{h}{J_{\hat{m}}N^{\hat{m}-1}} \right) + N - 1 \right] = \eta_{\hat{m}-1} \leq \sigma.$$

Observe that (3.28) is continuous in $\eta_{\hat{m}}$. Hence, if $\varphi_{\hat{m}} \in [\lceil \hat{s} \rceil(1-\epsilon), \lceil \hat{s} \rceil(1+\epsilon)]$ for some $\epsilon > 0$, and $\varphi_{\hat{m}-1}$ is equal to (3.28) with $\eta_{\hat{m}}$ replaced by $\varphi_{\hat{m}}$, then

$$(3.29) \quad \frac{1}{N} |\eta_{\hat{m}-1} - \varphi_{\hat{m}-1}| \leq \left(\frac{J_{\hat{m}+1}}{J_{\hat{m}}} \right) |\eta_{\hat{m}} - \varphi_{\hat{m}}| = \epsilon O(1) + \frac{2}{N}.$$

Since we already know from the reasoning leading up to Corollary 3.3 that any global maximum M must satisfy $a_i = 0$ for $i > \hat{m}$ and $a_{\hat{m}} \in [\lceil \hat{s} \rceil(1-\epsilon), \lceil \hat{s} \rceil(1+\epsilon)]$, by (3.29) we also have that $a_{\hat{m}-1} \in [\eta_{\hat{m}-1}(1-\epsilon'), \eta_{\hat{m}-1}(1+\epsilon')]$, with ϵ' allowed to be arbitrarily small as $N \rightarrow \infty$. We can now repeat these computations recursively, to conclude the same for $a_{\hat{m}-2}, \dots, a_0$.

Given $\eta_{\hat{m}}, \dots, \eta_{\hat{m}-i}$, let $0 \leq \sigma < N$ and $s(i, j) = \sum_{t=0}^i \eta_{\hat{m}-t} N^{\hat{m}-t} + j N^{\hat{m}-i-1}$, and note that

$$(3.30) \quad \begin{aligned} \mathcal{H}(\gamma_{s(i, \sigma)}) &= \mathcal{H}(\gamma_{s(i, 0)}) + J_{\hat{m}-i} N^{2(\hat{m}-i-1)} \sigma(N-\sigma) \\ & \quad + \sum_{j=1}^{i+1} J_{\hat{m}-i+j} N^{2(\hat{m}-i-1)+j} \sigma(N-2\eta_{\hat{m}-i+j-1}-1) \\ & \quad + \left(1 - \frac{1}{N}\right) \sum_{j=\hat{m}+2}^n \sigma J_j N^{\hat{m}-i-1+j} - h\sigma N^{\hat{m}-i-1}. \end{aligned}$$

Thus, we have

$$(3.31) \quad \begin{aligned} \mathcal{H}(\gamma_{s(i,\sigma+1)}) &= \mathcal{H}(\gamma_{s(i,\sigma)}) + J_{\hat{m}-i} N^{2(\hat{m}-i-1)} (N - 2\sigma - 1) \\ &\quad + \sum_{j=1}^{i+1} J_{\hat{m}-i+j} N^{2(\hat{m}-i-1)+j} (N - 2\eta_{\hat{m}-i+j-1} - 1) \\ &\quad + \left(1 - \frac{1}{N}\right) \sum_{j=\hat{m}+2}^n J_j N^{\hat{m}-i-1+j} - h N^{\hat{m}-i-1}, \end{aligned}$$

and hence $\mathcal{H}(\gamma_{s(i,\sigma+1)}) - \mathcal{H}(\gamma_{s(i,\sigma)}) \leq 0$ whenever

$$(3.32) \quad \begin{aligned} 0 \vee \left[\frac{1}{2} \left(\sum_{j=1}^{i+1} \left(\frac{J_{\hat{m}-i+j}}{J_{\hat{m}-i}} \right) N^j (N - 2\eta_{\hat{m}-i} - 1) \right. \right. \\ \left. \left. + \left(1 - \frac{1}{N}\right) \sum_{j=2}^{n-\hat{m}} \left(\frac{J_{\hat{m}+j}}{J_{\hat{m}-i}} \right) N^{i+j+1} - \frac{h}{J_{\hat{m}-i} N^{\hat{m}-i-1}} + N - 1 \right) \right] = \eta_{\hat{m}-i-1} \leq \sigma. \end{aligned}$$

Again it follows that if $\varphi_{\hat{m}-i} \in \{0, \dots, N-1\}$ and $\varphi_{\hat{m}-i-1}$ is equal to the left-hand side of (3.32) with $\eta_{\hat{m}-i}$ replaced by $\varphi_{\hat{m}-i}$ in (3.32), then

$$(3.33) \quad |\eta_{\hat{m}-i-1} - \varphi_{\hat{m}-i-1}| \leq \left(\frac{J_{\hat{m}-i+1}}{J_{\hat{m}-i}} \right) |\eta_{\hat{m}-i} - \varphi_{\hat{m}-i}| + \frac{2}{N}.$$

This proves the statement of the lemma. \square

We need to look at the change in energy when we go from a critical configuration in the set \mathcal{C}^* to a neighbouring configuration obtained by changing the sign at one vertex. Our next observation concerns the sets U_σ^- and U_σ^+ defined in the statement of Lemma 1.6.

Lemma 3.6. *Suppose that (A1) holds and that every $\xi \in \mathcal{C}^*$ has the same volume $|\xi| = k^*$, and that every configuration of volume k^* has energy at least $\Phi(\boxminus, \boxplus)$. Suppose furthermore that for every configuration $\sigma \in U_\xi^+$, $\mathcal{H}(\sigma) \neq \Phi(\boxminus, \boxplus)$. Then (1.22) is satisfied.*

Proof. Let $\xi \in \mathcal{C}^*$, and suppose that $\sigma \in U_\xi^-$, so that $\sigma = \xi \setminus \{v_a\}$ for some $a < k^*$. If σ lies on some optimal path, then, by the assumption that this path has a unique maximum, (1.22) is satisfied. Else, since ξ lies on an optimal path, there exists some configuration $\xi' = \xi \setminus \{v_b\}$ on the same path, of volume $|\xi'| = k^* - 1$ (note that by (A1) $k^* > 0$) and with $\Phi(\xi', \boxminus) < \Phi(\boxminus, \boxplus)$. We claim that the path $\sigma \rightarrow \sigma \cap \xi' \rightarrow \xi'$ stays strictly below $\Phi(\boxminus, \boxplus)$, which proves the statement of the lemma. Since by definition $\mathcal{H}(\sigma) < \Phi(\boxminus, \boxplus)$ and $\mathcal{H}(\xi') < \Phi(\boxminus, \boxplus)$, we only need to show that $\mathcal{H}(\sigma \cap \xi') < \Phi(\boxminus, \boxplus)$. However, note that

$$(3.34) \quad \begin{aligned} \mathcal{H}(\sigma \cap \xi') - \mathcal{H}(\sigma) &= \sum_{\substack{i \leq k^* \\ i \neq b, a}} J_{d(v_b, v_i)} - \sum_{i > k^*} J_{d(v_b, v_i)} - J_{d(v_b, v_a)} + h \\ &\leq \sum_{\substack{i \leq k^* \\ i \neq b}} J_{d(v_b, v_i)} - \sum_{i > k^*} J_{d(v_b, v_i)} + h = \mathcal{H}(\xi') - \mathcal{H}(\xi) < 0, \end{aligned}$$

where the last inequality uses the fact that $\Phi(\xi', \boxminus) < \Phi(\xi, \boxminus)$. This proves the claim for $\sigma \in U_\xi^-$. The argument for $\sigma \in U_\xi^+$ makes use of the fact that by assumption $\mathcal{H}(\sigma) \neq \Phi(\boxminus, \boxplus)$, and is otherwise identical to the argument above. \square

Next, let us first consider any configuration γ_k lying on the path γ , with $k = a_{n-1}N^{n-1} + \dots + a_0$, and let σ_b be a configuration obtained from γ_k by flipping the sign at a vertex w such that $d(w, v_k) =$

b for $b \in \{1, \dots, n\}$. Note that by symmetry it makes no difference which particular vertex we select. If $\sigma_b(w) = -\gamma_k(w) = -1$, then for $b = 1$ we have

$$\text{eq:vertexflip-s1} \quad (3.35) \quad \mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_k) = J_1(2a_0 - N - 1) + \sum_{i=1}^{n-1} J_{i+1}N^i(2a_i - N + 1) + h,$$

while for $2 \leq b \leq n$,

$$\text{eq:vertexflip-s2} \quad (3.36) \quad \begin{aligned} & \mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_k) \\ &= \sum_{i=1}^{b-1} J_i N^i \left(1 - \frac{1}{N}\right) + J_b \left(2 \sum_{i=0}^{b-1} a_i N^i - N^b - N^{b-1}\right) + \sum_{i=b}^{n-1} J_{i+1} N^i (2a_i - N + 1) + h. \end{aligned}$$

Similarly, if $\sigma_b(w) = -\gamma_k(w) = +1$, then for $b = 1$ we have

$$\text{eq:vertexflip+s1} \quad (3.37) \quad \mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_k) = \sum_{i=0}^{n-1} J_{i+1} N^i (N - 2a_i - 1) - h,$$

while for $2 \leq b \leq n$,

$$\text{eq:vertexflip+s2} \quad (3.38) \quad \begin{aligned} & \mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_k) \\ &= \sum_{i=1}^{b-1} J_i N^i \left(1 - \frac{1}{N}\right) + J_b \left(N^b - 2 \sum_{i=0}^{b-1} a_i N^i - N^{b-1}\right) + \sum_{i=b}^{n-1} J_{i+1} N^i (N - 2a_i - 1) - h. \end{aligned}$$

Under Assumption (A5), $\{\mathcal{H}(\gamma_i)\}_{i=0}^{N^n}$ attains a unique maximum. Indeed, this is immediate from (3.4). Furthermore, from Assumption (A4) it follows that \vec{J} is strictly monotone, and hence by Lemma 2.2 the path γ is strictly optimal. This implies that all $\sigma \in \mathcal{C}^*$ must have the same volume, and that every other configuration of that volume has larger energy. Hence the conditions of Lemma 3.6 are met.

3.4. Proof of Theorem 1.16. Define

$$\text{eq:Bd Bu} \quad (3.39) \quad \begin{aligned} B_d &= \left\{ 1 \leq b \leq \hat{m}: \sum_{i=1}^{b-1} J_i N^i \left(1 - \frac{1}{N}\right) + J_b \left(2 \sum_{i=0}^{b-1} \eta_i N^i - N^b - N^{b-1}\right) \right. \\ &\quad \left. + \sum_{i=b}^{n-1} J_{i+1} N^i (2\eta_i - N + 1) + h < 0 \right\} \\ B_u &= \left\{ 1 \leq b \leq n: \sum_{i=1}^{b-1} J_i N^i \left(1 - \frac{1}{N}\right) + J_b \left(N^b - 2 \sum_{i=0}^{b-1} \eta_i N^i - N^{b-1}\right) \right. \\ &\quad \left. + \sum_{i=b}^{n-1} J_{i+1} N^i (N - 2\eta_i - 1) - h < 0 \right\}, \end{aligned}$$

where $\{\eta_i\}_{i=0}^{n-1}$ is defined as in the statement of Lemma 3.5. By (3.36) and (3.38), B_d gives the distances to the ‘critical’ vertex of the vertices that are flipped in obtaining configurations that result in a lower energy than the critical configuration. Thus

$$(3.40) \quad \begin{aligned} N^-(\sigma) &= |\{\sigma \in U_\sigma^- : \mathcal{H}(\sigma) < \mathcal{H}(\sigma)\}| = [1 + o_N(1)] \sum_{i \in B_d} \eta_{i-1} N^{i-1}, \\ N^+(\sigma) &= |\{\sigma \in U_\sigma^+ : \mathcal{H}(\sigma) < \mathcal{H}(\sigma)\}| = [1 + o_N(1)] \sum_{i \in B_u} (N^i - \eta_{i-1} N^{i-1}). \end{aligned}$$

Hence, by Lemma 1.6, we have

$$\begin{aligned}
\frac{1}{K^*} &= [1 + o_N(1)] \sum_{\sigma \in \mathcal{C}^*} \frac{\left(\sum_{i \in B_d} \eta_{i-1} N^{i-1}\right) \left(\sum_{i \in B_u} (N^i - \eta_{i-1} N^{i-1})\right)}{\left(\sum_{i \in B_d} \eta_{i-1} N^{i-1}\right) + \left(\sum_{i \in B_u} (N^i - \eta_{i-1} N^{i-1})\right)} \\
(3.41) \quad &= [1 + o_N(1)] \frac{\left(\sum_{i \in B_d} \eta_{i-1} N^{i-1}\right) \left(\sum_{i \in B_u} (N^i - \eta_{i-1} N^{i-1})\right)}{\left(\sum_{i \in B_d} \eta_{i-1} N^{i-1}\right) + \left(\sum_{i \in B_u} (N^i - \eta_{i-1} N^{i-1})\right)} \\
&\quad \times \frac{N^{n-\hat{m}-1}}{N - \eta_0} \prod_{i=0}^{\hat{m}} \binom{N}{\eta_i} (N - \eta_i).
\end{aligned}$$

3.5. Proof of Theorems 1.13–1.14. Let $\{\tilde{J}_i\}_{i=1}^n$ be such that $\tilde{J}_i/N \rightarrow 0$ for all $i \in \{1, \dots, n\}$ as $N \rightarrow \infty$, and take $J_i = \tilde{J}_i/N^i$. It is easy to check that Assumption (A3) is satisfied given that Assumption (A2)(b) is also satisfied.

Proof of Theorem 1.13. From (1.32) and (3.32) we learn that

$$\begin{aligned}
\eta_{\hat{m}} &= \lceil \hat{s} \rceil = \left\lceil \frac{N}{2\tilde{J}_{\hat{m}+1}} \left(\left(1 - \frac{1}{N}\right) \sum_{i=\hat{m}+1}^n \tilde{J}_i - h \right) \right\rceil \\
(3.42) \quad &= [1 + o_N(1)] \frac{1}{2} \frac{N}{\tilde{J}_{\hat{m}+1}} \left(\sum_{i=\hat{m}+1}^n \tilde{J}_i - h \right)
\end{aligned}$$

and

$$\begin{aligned}
\eta_{\hat{m}-1} &= [1 + o_N(1)] \frac{N}{2}, \\
(3.43) \quad \eta_{\hat{m}-i} &= [1 + o_N(1)] \frac{N}{2} \left(\sum_{j=1}^{i+1} \left(\frac{\tilde{J}_{\hat{m}-i+j}}{\tilde{J}_{\hat{m}-i}} \right) \left(1 - \frac{2\eta_{\hat{m}-i+1}}{N}\right) + \sum_{j=2}^{n-\hat{m}} \left(\frac{\tilde{J}_{\hat{m}+j}}{\tilde{J}_{\hat{m}-i}} \right) - \frac{h}{\tilde{J}_{\hat{m}-i}} + 1 \right),
\end{aligned}$$

for $i = 1, \dots, \hat{m}$. This identifies the configurations announced in (1.46). \square

Proof of Theorem 1.14. Observe from (1.32) that

$$(3.44) \quad \hat{s} = \frac{N}{2\tilde{J}_{\hat{m}+1}} \left(\left(1 - \frac{1}{N}\right) \sum_{i=\hat{m}+1}^n \tilde{J}_i - h \right),$$

and by assumption (A1)(b) we have that

$$(3.45) \quad \lim_{N \rightarrow \infty} \left(1 - \frac{1}{N}\right) \sum_{i=\hat{m}+1}^n \tilde{J}_i - h = \lim_{N \rightarrow \infty} \sum_{i=\hat{m}+1}^n \tilde{J}_i - h > 0.$$

Then

$$\begin{aligned}
(3.46) \quad &\frac{k \sum_{i=\hat{m}+1}^n J_i N^i}{\lceil \hat{s} \rceil (2\hat{s} - \lceil \hat{s} \rceil + 1) J_{\hat{m}+1} N^{2\hat{m}}} = \frac{k \sum_{i=\hat{m}+1}^n \tilde{J}_i}{\lceil \hat{s} \rceil (2\hat{s} - \lceil \hat{s} \rceil + 1) \tilde{J}_{\hat{m}+1} N^{\hat{m}-1}} \\
&\leq \frac{N \sum_{i=\hat{m}+1}^n \tilde{J}_i}{\lceil \hat{s} \rceil (2\hat{s} - \lceil \hat{s} \rceil + 1) \tilde{J}_{\hat{m}+1}} = O(N^{-1}) \left(\tilde{J}_{\hat{m}+1} \right)^{-1} \sum_{i=\hat{m}+1}^n \tilde{J}_i,
\end{aligned}$$

and similarly

$$\begin{aligned} & \frac{\sum_{i=0}^{\hat{m}-1} J_{i+1} N^i \left((N - a_i - 1) \left(\sum_{j=0}^i a_j N^j \right) + a_i \left(N^i - \sum_{j=0}^{i-1} a_j N^j \right) \right)}{\lceil \hat{s} \rceil (2\hat{s} - \lceil \hat{s} \rceil + 1) J_{\hat{m}+1} N^{2\hat{m}}} \\ \text{eq: A2 check2} \quad (3.47) \quad & \leq \frac{N \sum_{i=0}^{\hat{m}-1} \tilde{J}_{i+1}}{\lceil \hat{s} \rceil (2\hat{s} - \lceil \hat{s} \rceil + 1) \tilde{J}_{\hat{m}+1}} = O(N^{-1}) \left(\tilde{J}_{\hat{m}+1} \right)^{-1} \sum_{i=\hat{m}+1}^n \tilde{J}_i. \end{aligned}$$

Summing (3.46) and (3.47) we get Assumption (A2). From (3.22) we get

$$(3.48) \quad \Gamma^* = [1 + o_N(1)] \frac{1}{4} N^{\hat{m}+1} \left(\tilde{J}_{\hat{m}+1} \right)^{-1} \left(\left(1 - \frac{1}{N} \right) \sum_{i=\hat{m}+1}^n \tilde{J}_i - h \right)^2.$$

□

4. STANDARD INTERACTION

In this section we consider the special case

$$\text{eq: regulardef} \quad (4.1) \quad J_i = \tilde{J}/N^i, \quad 1 \leq i \leq n,$$

for some $\tilde{J} > 0$. The Hamiltonian in (1.28) becomes

$$(4.2) \quad \mathcal{H}(h; \sigma) = -\frac{\tilde{J}}{2} \sum_{\substack{a, b \in \Lambda_N^n \\ a \neq b}} N^{-d(v_a, v_b)} \sigma(v_a) \sigma(v_b) - \frac{h}{2} \sum_{a \in \Lambda_N^n} \sigma(v_a),$$

where we exhibit the dependence on h . In Sections 4.1 we show that the energy landscape has certain symmetries. In Section 4.2 we exploit these symmetries to identify the location of the global maximum of the energy along the reference path γ . In Section 4.3 we use these results to prove Theorems 1.11 and 1.15. In Section 4.4 we compute the prefactor and prove Theorem 1.17.

4.1. Symmetries in the energy landscape. In this section we derive four lemmas (Lemmas 4.2–4.5 below) exhibiting certain symmetries in the energy landscape for the case of standard interaction (see Fig. 4.1). These symmetries will be crucial later on.

For any $h_1, h_2 > 0$ and $0 \leq a, b \leq N^n$,

$$\text{eq: differenthdifference} \quad (4.3) \quad \mathcal{H}(h_1; \gamma_a) - \mathcal{H}(h_1; \gamma_b) = \mathcal{H}(h_2; \gamma_a) - \mathcal{H}(h_2; \gamma_b) + (h_2 - h_1)(a - b).$$

Definition 4.1. A sequence $\{a_i\}_{i=1}^M \in \mathbb{R}^M$ is called symmetric when

$$(4.4) \quad a_i = a_{M-i+1}, \quad 1 \leq i \leq M,$$

and concave when

$$\text{eq: concavityproperty} \quad (4.5) \quad a_i - a_{i-1} \geq a_{i+1} - a_i, \quad 2 \leq i \leq M - 1.$$

The following lemma is elementary.

Lemma 4.2. Suppose that the sequence $\{a_i\}_{i=1}^M$ is symmetric and concave. Then

$$(4.6) \quad \max_{1 \leq i \leq M} a_i = a_{\lceil \frac{M}{2} \rceil}.$$

Recall the definition of \hat{m} from (1.31), and note that now

$$\text{eq: mhatregular} \quad (4.7) \quad \hat{m}_h = \left\lceil n - \frac{h}{\tilde{J}} \left(1 - \frac{1}{N} \right)^{-1} \right\rceil,$$

where again we exhibit the dependence on h . It was shown in Section 3.2 that, in the hierarchical limit $N \rightarrow \infty$, \hat{m}_h gives the order of magnitude of a critical configuration (in particular, the asymptotic size of a critical configuration was shown to be $\hat{s}N^{\hat{m}}$). We will now show that for the standard interaction in (4.1), \hat{m}_h plays a similar role.

Let $\gamma: \Xi \rightarrow \Theta$ be the optimal path defined in Section 1.4. We begin by considering the Hamiltonian $i \mapsto \mathcal{H}(h; \gamma_i)$ for certain special values of h . Recall $h^{(m,s)}$ defined in (1.39). In terms of this quantity, we have

$$\begin{aligned}
(4.8) \quad & \mathcal{H}\left(h^{(m,s)}; \gamma_{sN^{\hat{m}}}\right) - \mathcal{H}\left(h^{(m,s)}; \Xi\right) \\
&= \frac{\tilde{J}}{N} s N^m (N-s) + \tilde{J} s N^m \sum_{i=m+2}^n \left(1 - \frac{1}{N}\right) - s h^{(m,s)} N^m \\
&= s N^m \left(\tilde{J} (N-s) \frac{1}{N} + \tilde{J} \left(1 - \frac{1}{N}\right) (n-m-1) - h^{(m,s)} \right) = 0
\end{aligned}$$

and

$$\text{eq:mhat-hms} \quad (4.9) \quad \hat{m}_{h^{(m,s)}} = \left\lfloor m + (s-1) \frac{1}{N} \left(1 - \frac{1}{N}\right)^{-1} \right\rfloor = m.$$

A magnetic field that takes the form $h^{(m,s)}$ gives rise to symmetries in the energy landscape along the path γ , which we can exploit in order to find the values at which $i \mapsto \mathcal{H}(h^{(m,s)}; \gamma_i)$ attains its global maximum. Later we will use this information to find the location of the global maxima for general values of h . First we show that the global maximum of $i \mapsto \mathcal{H}(h^{(m,s)}; \gamma_i)$ is attained in the interval $[0, sN^m]$.

Lemma 4.3. *For any $1 \leq s \leq N$ and $0 \leq m \leq n-1$,*

$$\text{eq:Hineq} \quad (4.10) \quad \max_{1 \leq i \leq N^n} \mathcal{H}\left(h^{(m,s)}; \gamma_i\right) = \max_{i \leq sN^m} \mathcal{H}\left(h^{(m,s)}; \gamma_i\right).$$

Proof. Let $K = a_{n-1}N^{n-1} + \dots + a_0$ and $u(i) = a_{n-1}N^{n-1} + \dots + a_i N^i$, and note that, by Lemma 3.2,

$$(4.11) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_{u(m+1)}\right) \leq \mathcal{H}\left(h^{(m,s)}; \gamma_{u(m+2)}\right) + \mathcal{H}\left(h^{(m,s)}; \gamma_{a_{m+1}N^{m+1}}\right) - \mathcal{H}\left(h^{(m,s)}; \Xi\right).$$

By Lemma 3.1 and the definition of $m = \hat{m}$ in (3.12), we have, for $0 \leq m < n-1$,

$$\begin{aligned}
(4.12) \quad & \mathcal{H}\left(h^{(m,s)}; \gamma_{a_{m+1}N^{m+1}}\right) - \mathcal{H}\left(h^{(m,s)}; \gamma_{(a_{m+1}-1)N^{m+1}}\right) \\
& \leq \mathcal{H}\left(h^{(m,s)}; \gamma_{N^{m+1}}\right) - \mathcal{H}\left(h^{(m,s)}; \Xi\right) \leq 0.
\end{aligned}$$

Hence, by induction,

$$\begin{aligned}
\text{eq:inductsetp} \quad (4.13) \quad & \mathcal{H}\left(h^{(m,s)}; \gamma_{a_{m+1}N^{m+1}}\right) \leq \mathcal{H}\left(h^{(m,s)}; \gamma_{(a_{m+1}-1)N^{m+1}}\right) \\
& \leq \dots \leq \mathcal{H}\left(h^{(m,s)}; \gamma_{N^{m+1}}\right) \leq \mathcal{H}\left(h^{(m,s)}; \Xi\right)
\end{aligned}$$

and therefore

$$(4.14) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_{u(m+1)}\right) \leq \mathcal{H}\left(h^{(m,s)}; \gamma_{u(m+2)}\right).$$

Once again it follows from inductive reasoning that

$$(4.15) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_{u(m+1)}\right) \leq \mathcal{H}\left(h^{(m,s)}; \gamma_{a_{n-1}N^{n-1}}\right).$$

By the same reasoning as in (4.13), we have

$$(4.16) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_{a_{n-1}N^{n-1}}\right) \leq \mathcal{H}\left(h^{(m,s)}; \gamma_{N^{n-1}}\right) \leq \mathcal{H}\left(h^{(m,s)}; \Xi\right)$$

and hence

$$(4.17) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_{u(m+1)}\right) \leq \mathcal{H}\left(h^{(m,s)}; \Xi\right).$$

Thus

$$\begin{aligned}
& \mathcal{H}\left(h^{(m,s)}; \gamma_K\right) - \mathcal{H}\left(h^{(m,s)}; \boxplus\right) \\
\text{eq:slowing2-inequality} \quad (4.18) \quad &= \mathcal{H}\left(h^{(m,s)}; \gamma_K\right) - \mathcal{H}\left(h^{(m,s)}; \gamma_{u(m+1)}\right) + \mathcal{H}\left(h^{(m,s)}; \gamma_{u(m+1)}\right) - \mathcal{H}\left(h^{(m,s)}; \boxplus\right) \\
&\leq \mathcal{H}\left(h^{(m,s)}; \gamma_K\right) - \mathcal{H}\left(h^{(m,s)}; \gamma_{u(m+1)}\right) \leq \mathcal{H}\left(h^{(m,s)}; \gamma_{a_m N^m + \dots + a_0}\right) - \mathcal{H}\left(h^{(m,s)}; \boxplus\right),
\end{aligned}$$

where the last inequality again follows from Lemma 3.2. Moreover, for $m = n - 1$ the inequality in (4.18) is immediate. If $a_m < s$, then the claim in (4.10) follows immediately. Otherwise we have $\mathcal{H}\left(h^{(m,s)}; \gamma_{a_m N^m}\right) \leq \mathcal{H}\left(h^{(m,s)}; \boxplus\right)$ and hence, by Lemma 3.2 and using the abbreviation $v(i) = a_i N^i + \dots + a_0$,

$$\begin{aligned}
& \mathcal{H}\left(h^{(m,s)}; \gamma_{v(m)}\right) - \mathcal{H}\left(h^{(m,s)}; \boxplus\right) \\
(4.19) \quad &\leq \mathcal{H}\left(h^{(m,s)}; \gamma_{v(m)}\right) - \mathcal{H}\left(h^{(m,s)}; \gamma_{a_m N^m}\right) + \mathcal{H}\left(h^{(m,s)}; \gamma_{a_m N^m}\right) - \mathcal{H}\left(h^{(m,s)}; \boxplus\right) \\
&\leq \mathcal{H}\left(h^{(m,s)}; \gamma_{v(m-1)}\right) - \mathcal{H}\left(h^{(m,s)}; \boxplus\right) \\
&\leq \max_{1 \leq i \leq s N^m} \mathcal{H}\left(h^{(m,s)}; \gamma_i\right) - \mathcal{H}\left(h^{(m,s)}; \boxplus\right),
\end{aligned}$$

which settles the claim. \square

We next derive two results stating $\{\mathcal{H}(h^{(m,s)}, \gamma_i)\}_{i=1}^{s N^m}$ (illustrated in Fig. 4.1) is symmetric and fractal-like, which is used later to locate the global maxima of this sequence.

Lemma 4.4. *The sequence $\{\mathcal{H}(h^{(m,s)}; \gamma_i)\}_{i=0}^{s N^m}$ is symmetric, i.e.,*

$$(4.20) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_K\right) = \mathcal{H}\left(h^{(m,s)}; \gamma_{s N^m - K}\right), \quad 0 \leq K \leq s N^m.$$

Proof. Let $K = k_{n-1} N^{n-1} + \dots + k_0$, so that

$$\begin{aligned}
& \mathcal{H}\left(h^{(m,s)}; \gamma_K\right) - \mathcal{H}\left(h^{(m,s)}; \boxplus\right) + h^{(m,s)} K \\
(4.21) \quad &= \sum_{i=0}^{n-1} J_{i+1} N^i \left(\left(\sum_{j=0}^i k_j N^j \right) (N - k_i - 1) + k_i \left(N^i - \sum_{j=0}^{i-1} k_j N^j \right) \right) \\
&= \sum_{i=0}^{n-1} J_{i+1} N^i \left(\left(\sum_{j=0}^{i-1} k_j N^j \right) (N - k_i - 1) + k_i N^i (N - k_i - 1) + k_i N^i - k_i \sum_{j=0}^{i-1} k_j N^j \right) \\
&= \sum_{i=0}^{n-1} J_{i+1} N^i \left(\left(\sum_{j=0}^{i-1} k_j N^j \right) (N - 2k_i - 1) + k_i N^i (N - k_i) \right) \\
&= \sum_{i=0}^{n-1} \frac{\tilde{J}}{N} \left(\left(\sum_{j=0}^{i-1} k_j N^j \right) (N - 2k_i - 1) + k_i N^i (N - k_i) \right).
\end{aligned}$$

Since $k_i = 0$ for $i > m$ and $k_m < s$, this simplifies to

$$\begin{aligned}
& \mathcal{H}\left(h^{(m,s)}; \gamma_K\right) - \mathcal{H}\left(h^{(m,s)}; \boxplus\right) \\
\text{eq:HKregsymmetric} \quad (4.22) \quad &= \sum_{i=0}^m \frac{\tilde{J}}{N} \left(\left(\sum_{j=0}^{i-1} k_j N^j \right) (N - 2k_i - 1) + k_i N^i (N - k_i) \right) \\
&\quad + K \left(\tilde{J} \left(1 - \frac{1}{N} \right) (n - m - 1) - h^{(m,s)} \right).
\end{aligned}$$

Note that if $\tilde{K} = sN^m - K$, then the number of interacting pairs at distance $i = 0, \dots, m$ in the configuration $\gamma_{\tilde{K}}$ (i.e., vertices v_a, v_b such that $\gamma_{\tilde{K}}(v_a) = -\gamma_{\tilde{K}}(v_b)$ and $d(v_a, v_b) = i$) is the same as in the configuration γ_K . At distance $m+1$ this number is equal to

$$(4.23) \quad N^m \left(K(s - k_m - 1) + \left(N^m - \sum_{j=0}^{m-1} k_j N^j \right) k_m + (sN^m - K)(N - s) \right)$$

and therefore we conclude that

$$(4.24) \quad \begin{aligned} & \mathcal{H}(h^{(m,s)}; \gamma_{\tilde{K}}) - \mathcal{H}(h^{(m,s)}; \boxplus) \\ &= \sum_{i=0}^{m-1} \frac{\tilde{J}}{N} \left(\left(\sum_{j=0}^{i-1} k_j N^j \right) (N - 2k_i - 1) + k_i N^i (N - k_i) \right) \\ &+ \frac{\tilde{J}}{N} \left(K(s - k_m - 1) + \left(N^m - \sum_{j=0}^{m-1} k_j N^j \right) k_m + (sN^m - K)(N - s) \right) \\ &+ \sum_{i=m+1}^{n-1} \tilde{J} \left(1 - \frac{1}{N} \right) \left(sN^m - \sum_{j=0}^m k_j N^j \right) - h^{(m,s)} \tilde{K}. \end{aligned}$$

Thus, we have

$$(4.25) \quad \begin{aligned} & \mathcal{H}(h^{(m,s)}; \gamma_{\tilde{K}}) - \mathcal{H}(h^{(m,s)}; \gamma_K) = \sum_{i=m+1}^{n-1} \tilde{J} \left(1 - \frac{1}{N} \right) (sN^m - 2K) \\ &+ \frac{\tilde{J}}{N} (K(s - k_m - 1) + (sN^m - K)(N - s) - K(N - k_i - 1)) - h^{(m,s)}(sN^m - 2K), \end{aligned}$$

which is equal to 0 if and only if

$$(4.26) \quad \begin{aligned} & h^{(m,s)}(sN^m - 2K) \\ &= \tilde{J} \left(1 - \frac{1}{N} \right) (sN^m - 2K)(n - m - 1) \\ &+ \frac{\tilde{J}}{N} (K(s - k_m - 1) + (sN^m - K)(N - s) - K(N - k_m - 1)) \\ &= \tilde{J} \left(1 - \frac{1}{N} \right) (sN^m - 2K)(n - m - 1) + \frac{\tilde{J}}{N} (K(s - N) + (sN^m - K)(N - s)) \\ &= \tilde{J} (sN^m - 2K) \left(\left(1 - \frac{1}{N} \right) (n - m) - (s - 1) \frac{1}{N} \right), \end{aligned}$$

which indeed is true by the definition of $h^{(m,s)}$ in (1.39). \square

To state the second result we need some more notation. Let $Q: \mathbb{N}_0 \rightarrow \{0, 1\}$ be defined by

$$(4.27) \quad \text{eq: def Q} \quad Q(a) = a \bmod 2.$$

For all integers $k \in \{1, \dots, m\}$ taking the form $k = a(1 + Q(N + 1)) - Q((N + 1)(s + 1))$ for some $a \in \{1, \dots, m\}$, define the integer intervals $S_k = [S_k^-, S_k^+]$, where

$$(4.28) \quad \text{eq: SetSk} \quad \begin{aligned} S_k^- &= \left(\left\lfloor \frac{s}{2} \right\rfloor - 1 + Q(s(N + 1)) \right) N^m + \sum_{j=1}^{k-1} a_{m-j} N^{m-j} + (1 + Q(sN)) N^{m-k}, \\ S_k^+ &= \left(\left\lfloor \frac{s}{2} \right\rfloor - 1 + Q(s(N + 1)) \right) N^m + \sum_{j=1}^{k-1} a_{m-j} N^{m-j} + N^{m-k+1}, \end{aligned}$$

and

$$(4.29) \quad a_{m-j} = \left\lfloor \frac{N}{2} - Q((j+s+1)(N+1)) \right\rfloor.$$

The following clarification regarding (4.28) is in order. For odd values of N , (4.28) defines the sets S_1, \dots, S_m , and the coefficients a_{m-j} are all equal to $\lfloor \frac{N}{2} \rfloor = \frac{N-1}{2}$. For even values of N and even values of s , (4.28) defines the odd-indexed sets $S_1, S_3, \dots, S_{2\lfloor \frac{m}{2} \rfloor + 1}$ and the coefficients a_{m-j} are given by $a_{m-1} = \frac{N}{2}$, $a_{m-2} = \frac{N}{2} - 1$, etc. For even values of N and odd values of s , (4.28) defines the even-indexed sets $S_2, S_4, \dots, S_{2\lfloor \frac{m}{2} \rfloor}$ and the coefficients a_{m-j} are given by $a_{m-1} = \frac{N}{2} - 1$, $a_{m-2} = \frac{N}{2}$, etc.

Lemma 4.5. *For every $k \in \{1, \dots, m\}$ that takes the form*

$$(4.30) \quad k = a(1 + Q(N+1)) + Q((N+1)(s+1))$$

for some $a \in \mathbb{N}_0$, the sequence $\{\mathcal{H}(h^{(m,s)}; \gamma_i)\}_{i \in S_k}$ is symmetric.

Proof. Suppose that $K \in S_k$, so that

$$(4.31) \quad K = \sum_{i=0}^m a_i N^i = \left(\left\lfloor \frac{s}{2} \right\rfloor - 1 + Q(s(N+1)) \right) N^m + \sum_{j=1}^{k-1} a_{m-j} N^{m-j} + R,$$

where

$$(4.32) \quad R = a_{m-k} N^{m-k} + a_{m-k-1} N^{m-k-1} + \dots + a_0$$

for $1 + Q(sN) \leq a_{m-k} \leq N - 1$ and $0 \leq a_i \leq N - 1$ for $0 \leq i < m - k$. Also let

$$(4.33) \quad \begin{aligned} \tilde{K} &= \left(\left\lfloor \frac{s}{2} \right\rfloor - 1 + Q(s(N+1)) \right) N^m \\ &+ \sum_{j=1}^{k-1} a_{m-j} N^{m-j} + N^{m-k+1} - R + (1 + Q(sN)) N^{m-k} \\ &= K + N^{m-k+1} - 2R + (1 + Q(sN)) N^{m-k}, \end{aligned}$$

so that K and \tilde{K} are mirrored points in S_k (i.e., if K is the i^{th} point in S_k , then \tilde{K} is the $(|S_k| - i)^{\text{th}}$ point). Note that, by (4.22),

$$(4.34) \quad \begin{aligned} \mathcal{H}(h^{(m,s)}; \gamma_K) - \mathcal{H}(h^{(m,s)}; \Xi) &= \sum_{i=0}^m \frac{\tilde{J}}{N} \left(\left(\sum_{j=0}^{i-1} a_j N^j \right) (N - 2a_i - 1) + a_i N^i (N - a_i) \right) \\ &+ K \left(\tilde{J} \left(1 - \frac{1}{N} \right) (n - m - 1) - h^{(m,s)} \right). \end{aligned}$$

Observe that, for $1 \leq i \leq m - k$, the total number of interacting pairs at distance i in $\gamma_{\tilde{K}}$ (i.e., vertices v, w such that $d(v, w) = i$ and $\gamma_{\tilde{K}}(v) = -\gamma_{\tilde{K}}(w)$), is the same as in γ_K . At distance $m - k + 1$, the number of interacting pairs in $\gamma_{\tilde{K}}$ is equal to the number of interacting pairs in γ_K plus $(1 + Q(sN))N^{m-k}(R - (1 + Q(sN))N^{m-k})$ minus $(1 + Q(sN))N^{m-k}(N^{m-k+1} - R)$. For $m - k + 2 \leq i$, the number of interacting pairs at distance i in $\gamma_{\tilde{K}}$ is equal to the number of interacting pairs in γ_K plus $a_i N^i (R - (1 + Q(sN))N^{m-k})$ minus $a_i N^i (N^{m-k+1} - R)$, and plus

$(N - a_i - 1)N^i(N^{m-k+1} - R)$ minus $(N - a_i - 1)N^i(R - (1 + Q(sN))N^{m-k})$. Thus, we have

$$\begin{aligned}
& \left(\mathcal{H}(h^{(m,s)}; \gamma_{\tilde{K}}) - \mathcal{H}(h^{(m,s)}; \Xi) \right) \left(\frac{\tilde{J}}{N} \right)^{-1} \\
&= \sum_{i=0}^{m-k} \left(\left(\sum_{j=0}^{i-1} a_j N^j \right) (N - 2a_i - 1) + a_i N^i (N - a_i) \right) \\
&\quad + (1 + Q(sN)) (2R - N^{m-k+1} - (1 + Q(sN)) N^{m-k}) \\
(4.35) \quad &+ \sum_{i=m-k+1}^m \left(\left(\sum_{j=0}^{i-1} a_j N^j \right) (N - 2a_i - 1) + a_i N^i (N - a_i) \right) \\
&\quad + \sum_{i=m-k+1}^m (N - 2a_i - 1) (N^{m-k+1} - 2R + (1 + Q(sN)) N^{m-k}) \\
&\quad + \sum_{i=m+1}^{n-1} \tilde{J} \left(1 - \frac{1}{N} \right) \left(\sum_{j=0}^m a_j N^j + (N^{m-k+1} - 2R + (1 + Q(sN)) N^{m-k}) \right) \\
&\quad - h^{(m,s)} \tilde{K}.
\end{aligned}$$

Hence it follows that
eq:HKtildeKdifference (4.36)

$$\begin{aligned}
& \mathcal{H}(h^{(m,s)}; \gamma_{\tilde{K}}) - \mathcal{H}(h^{(m,s)}; \gamma_K) = \frac{\tilde{J}}{N} (1 + Q(sN)) (2R - N^{m-k+1} - (1 + Q(sN)) N^{m-k}) \\
&\quad + \sum_{i=m-k+1}^m \frac{\tilde{J}}{N} (N - 2a_i - 1) (N^{m-k+1} - 2R + (1 + Q(sN)) N^{m-k}) \\
&\quad + \sum_{i=m+1}^{n-1} \tilde{J} \left(1 - \frac{1}{N} \right) (N^{m-k+1} - 2R + (1 + Q(sN)) N^{m-k}) - h^{(m,s)} (\tilde{K} - K).
\end{aligned}$$

Note that (4.36) is equal to zero if and only if

$$\begin{aligned}
& h^{(m,s)} (\tilde{K} - K) = h^{(m,s)} (N^{m-k+1} - 2R + N^{m-k}) \\
&= \frac{\tilde{J}}{N} (1 + Q(sN)) (2R - N^{m-k+1} - (1 + Q(sN)) N^{m-k}) \\
(4.37) \quad &+ \sum_{i=m-k+1}^m \frac{\tilde{J}}{N} (N - 2a_i - 1) (N^{m-k+1} - 2R + (1 + Q(sN)) N^{m-k}) \\
&\quad + \sum_{i=m+1}^{n-1} \tilde{J} \left(1 - \frac{1}{N} \right) (N^{m-k+1} - 2R + (1 + Q(sN)) N^{m-k}),
\end{aligned}$$

which holds whenever

$$\begin{aligned}
(4.38) \quad & h^{(m,s)} = -\frac{\tilde{J}}{N} (1 + Q(sN)) + \sum_{i=m-k+1}^m \frac{\tilde{J}}{N} (N - 2a_i - 1) + \sum_{i=m+1}^{n-1} \tilde{J} \left(1 - \frac{1}{N} \right) \\
&= -\frac{\tilde{J}}{N} (1 + Q(sN)) + \frac{\tilde{J}}{N} \sum_{i=m-k+1}^{m-1} (N - 2a_i - 1) \\
&\quad + \frac{\tilde{J}}{N} \left(N - 2 \left\lfloor \frac{s}{2} \right\rfloor + 1 - Q(s(N+1)) \right) + (n - m - 1) \tilde{J} \left(1 - \frac{1}{N} \right).
\end{aligned}$$

If N is odd, then $(N - 2a_i - 1) = (N - 2\lfloor \frac{N}{2} \rfloor - 1) = 0$, and hence $\sum_{i=m-k+1}^{m-1} (N - 2a_i - 1) = 0$. If N is even, then the terms $(N - 2a_i - 1)$ alternate between -1 and 1 . Thus, if s is even, k is odd and $\sum_{i=m-k+1}^{m-1} (N - 2a_i - 1) = 0$ because the sum has an even number of terms, while if s is odd, then the sum adds up to $\frac{\tilde{J}}{N}$. We can encode this as

$$(4.39) \quad \frac{\tilde{J}}{N} \sum_{i=m-k+1}^{m-1} (N - 2a_i - 1) = \frac{\tilde{J}}{N} Q(s(N+1)).$$

Recalling (1.39), it remains to show that

$$(4.40) \quad \begin{aligned} \left(1 - \frac{1}{N}\right)(n-m) - (s-1)\frac{1}{N} &= -\frac{1}{N}(1 + Q(sN)) + \frac{1}{N}Q(s(N+1)) \\ &+ \frac{1}{N}\left(N - 2\left\lfloor \frac{s}{2} \right\rfloor + 1 - 2Q(s(N+1))\right) + (n-m-1)\left(1 - \frac{1}{N}\right), \end{aligned}$$

or equivalently

$$(4.41) \quad -(s-1)\frac{1}{N} = -\frac{1}{N}\left(2\left\lfloor \frac{s}{2} \right\rfloor + Q(sN) - 1 + Q(s(N+1))\right) = -(s-1)\frac{1}{N},$$

which is trivially true. \square

The symmetries in Lemmas 4.2–4.5 are depicted in Fig. 4.1.




FIGURE 8. Plot of $i \mapsto \mathcal{H}(\gamma_i)$ for Λ_5^9 , with $\tilde{J} = 10.3$ and $h = h^{(m,s)} = \tilde{J}\left(1 - \frac{1}{N}\right)(n-m) - \frac{(s-1)}{N}$ with $m = 4$ and $s = 8$. The solid-line in the left plot corresponds to values $i = 0, 1, \dots, sN^m$, and is symmetric as shown in Lemma 4.4. The solid-line in the right plot shows symmetry of $\mathcal{H}(\gamma_i)$ for values $i \in S_1$, as shown in Lemma 4.5.

4.2. Global maximum along the reference path. In this section we derive two propositions (Propositions 4.6–4.7 below) identifying the location of the global maximum of $i \mapsto \mathcal{H}(h^{(m,s)}; \gamma_i)$.

Proposition 4.6. *Suppose that N is odd. If s is odd, then*

$$(4.42) \quad \begin{aligned} \text{eq:maxloc} \quad \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor sN^m/2 \rfloor}\right) &= \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor sN^m/2 \rfloor + 1}\right) \\ &= \max_{1 \leq i \leq sN^m} \mathcal{H}\left(h^{(m,s)}; \gamma_i\right) = \max_{1 \leq i \leq N^n} \mathcal{H}\left(h^{(m,s)}; \gamma_i\right), \end{aligned}$$

and for all $i < \lfloor sN^m/2 \rfloor$,

$$(4.43) \quad \text{eq:1st max s odd} \quad \mathcal{H}\left(h^{(m,s)}; \gamma_i\right) < \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor sN^m/2 \rfloor}\right).$$

If s is even, then

$$(4.44) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}\right) = \max_{1 \leq i \leq sN^m} \mathcal{H}\left(h^{(m,s)}; \gamma_i\right) = \max_{1 \leq i \leq N^n} \mathcal{H}\left(h^{(m,s)}; \gamma_i\right)$$

and for all $i < \lfloor (s-1)N^m/2 \rfloor + 1$,

$$(4.45) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_i\right) < \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}\right).$$

Proof. The first equality in (4.42) is immediate from Lemma 4.4 since $\lfloor sN^m/2 \rfloor + 1 = sN^m - \lfloor sN^m/2 \rfloor$, while the third equality follows from Lemma 4.3. We claim that the second equality in (4.42) follows from both Lemma 4.4 and Lemma 4.2. Indeed, note that by Lemma 3.1 the sequence

$$(4.46) \quad \{\mathcal{H}(h; \gamma_i)\}, \quad \left\lfloor \frac{sN^m}{2} \right\rfloor - \left\lfloor \frac{N}{2} \right\rfloor + 1 \leq i \leq \left\lfloor \frac{sN^m}{2} \right\rfloor + \left\lfloor \frac{N}{2} \right\rfloor + 2.$$

is concave, and by Lemma 4.4 is also symmetric. Therefore, by Lemma 4.2, we have that $\mathcal{H}(\gamma_i) \leq \mathcal{H}(\gamma_{\lfloor sN^m/2 \rfloor})$ for all i such that $d(v_i, v_{\lfloor sN^m/2 \rfloor}) = d(v_i, v_{\lfloor sN^m/2 \rfloor + 1}) = 1$. In fact, from Lemma 3.1 we have a strict form of concavity,

$$(4.47) \quad \begin{aligned} & \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor sN^m/2 \rfloor}\right) - \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor sN^m/2 \rfloor}\right) \\ &= \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor sN^m/2 \rfloor + 1}\right) - \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor sN^m/2 \rfloor}\right) + 2\tilde{J} = 2\tilde{J}, \end{aligned}$$

which shows that

$$(4.48) \quad \mathcal{H}(\gamma_{\lfloor sN^m/2 \rfloor}) > \mathcal{H}(\gamma_i) \quad \forall i < \lfloor sN^m/2 \rfloor : d(v_i, v_{\lfloor sN^m/2 \rfloor}) = d(v_i, v_{\lfloor sN^m/2 \rfloor + 1}) = 1.$$

Suppose that this is also true for all i such that $d(v_i, v_{\lfloor sN^m/2 \rfloor}) = r$, and let z be such that $d(v_z, v_{\lfloor sN^m/2 \rfloor}) = r + 1$. Note that if $r + 1 < m + 1$, then z belongs to a sequence of the form $\{z_0 + tN^r\}_{t=0}^{N-1}$ for some z_0 such that all N terms in the sequence belong to the same $(r + 1)$ -block, while if $r + 1 = m + 1$, then $z \in \{z_0 + tN^r\}_{t=0}^{s-1}$ such that again all s terms belong to the first $(m + 1)$ -block. Observe that the sequence $\{\mathcal{H}(h^{(m,s)}; \gamma_i)\}_{i \in A}$ is concave by Lemma 3.1 and symmetric by Lemma 4.4, where

$$(4.49) \quad A = \left\{ \{z_0 + tN^r\}_{t=0}^{N-1} \cap [0, \lfloor sN^m/2 \rfloor], \{sN^m - z_0 - tN^r\}_{t=0}^{N-1} \cap [\lfloor sN^m/2 \rfloor + 1, sN^m] \right\}$$

if $r + 1 < m + 1$, and

$$(4.50) \quad A = \left\{ \{z_0 + tN^r\}_{t=0}^{s-1} \cap [0, \lfloor sN^m/2 \rfloor], \{sN^m - z_0 - tN^r\}_{t=0}^{s-1} \cap [\lfloor sN^m/2 \rfloor + 1, sN^m] \right\}$$

if $r + 1 = m + 1$. Hence it attains its maximum only at the two midpoints of the sequence A (which has $N + 1$ terms in total). At least one of these two points is at distance r from $v_{\lfloor sN^m/2 \rfloor}$. Thus, by the inductive hypothesis we have that $\mathcal{H}(h^{(m,s)}; \gamma_z) < \mathcal{H}(h^{(m,s)}; \gamma_{\lfloor sN^m/2 \rfloor})$.

Next, we look at the case when s is even. By (4.3) and the above result for the odd value $s - 1$, we have that, for $t < \lfloor (s-1)N^m/2 \rfloor + 1$,

$$(4.51) \quad \begin{aligned} & \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}\right) - \mathcal{H}\left(h^{(m,s)}; \gamma_t\right) \\ & \geq \mathcal{H}\left(h^{(m,s-1)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}\right) - \mathcal{H}\left(h^{(m,s-1)}; \gamma_t\right) > 0, \end{aligned}$$

and thus we only need to show that

$$(4.52) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}\right) \geq \mathcal{H}\left(h^{(m,s)}; \gamma_t\right) \quad \forall \lfloor (s-1)N^m/2 \rfloor + 1 \leq t \leq \lfloor sN^m/2 \rfloor + 1.$$

To do this, recall first that by Lemma 4.5 the sequence $\{\mathcal{H}(h^{(m,s)}; \gamma_i)\}_{i \in S_m}$ is symmetric, concave and of odd cardinality. Furthermore, $\mathcal{H}(h^{(m,s)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1})$ is the midpoint of this sequence, and for all $i, j \in S_m$ we have $d(v_i, v_j) = 1$. Hence

$$(4.53) \quad \mathcal{H}\left(h^{(m,s)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}\right) > \mathcal{H}\left(h^{(m,s)}; \gamma_i\right)$$

for all $i < \lfloor (s-1)N^m/2 \rfloor + 1$ such that $d(v_i, v_{\lfloor (s-1)N^m/2 \rfloor + 1}) = 1$. Now observe that $S_m \subset S_{m-1} \subset \dots \subset S_1$, and suppose that $\mathcal{H}(h^{(m,s)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}) > \mathcal{H}(h^{(m,s)}; \gamma_i)$ for all $i < \lfloor (s-1)N^m/2 \rfloor + 1$ such that $d(v_i, v_{\lfloor (s-1)N^m/2 \rfloor + 1}) = r$. If i is such that $d(v_i, v_{\lfloor (s-1)N^m/2 \rfloor + 1}) = r + 1$, then like in the s -odd case we can construct a concave and symmetric sequence such that the midpoint (and hence maximum) of this sequence is at distance r or less from $v_{\lfloor (s-1)N^m/2 \rfloor + 1}$. It follows that (4.44) and (4.45) hold. \square

Proposition 4.7. *Suppose that N is even, and let*

$$\text{eq:max-index-r} \quad (4.54) \quad r = \left(\frac{s-1}{2} \right) N^m + \sum_{j=1}^{m-1} a_{m-j} N^{m-j} + \frac{N}{2},$$

where

$$\text{eq:amj-Neven} \quad (4.55) \quad a_{m-j} = \frac{N}{2} - Q(j+s+1).$$

If $s = 2a + 1$ for some $a \in \{0, \dots, \frac{N}{2} - 1\}$, then

$$\text{eq:maxloc-1} \quad (4.56) \quad \mathcal{H}(h^{(m,s)}; \gamma_r) = \max_{1 \leq i \leq sN^m} \mathcal{H}(h^{(m,s)}; \gamma_i) = \max_{1 \leq i \leq N^n} \mathcal{H}(h^{(m,s)}; \gamma_i)$$

and, for all $i < r$,

$$(4.57) \quad \mathcal{H}(h^{(m,s)}; \gamma_i) < \mathcal{H}(h^{(m,s)}; \gamma_r).$$

Similarly,

$$\text{eq:maxloc2-1} \quad (4.58) \quad \mathcal{H}(h^{(m,s+1)}; \gamma_r) = \max_{1 \leq i \leq (s+1)N^m} \mathcal{H}(h^{(m,s+1)}; \gamma_i) = \max_{1 \leq i \leq N^n} \mathcal{H}(h^{(m,s+1)}; \gamma_i)$$

and, for all $i < r$,

$$(4.59) \quad \mathcal{H}(h^{(m,s+1)}; \gamma_i) < \mathcal{H}(h^{(m,s)}; \gamma_r).$$

Proof. The coordinates a_{m-j} are defined below (4.28). Noting that r is the midpoint of the smallest of the sets $\{S_k\}$ (for odd or even indices k depending on the case in question), we see that the claim follows from similar computations as those in the proof of Proposition 4.6. \square

4.3. Proof of Theorems 1.11 and 1.15. We now use Propositions 4.6 and 4.7 to determine the size of the critical configurations and prove Theorems 1.15 and 1.11 (Propositions 4.8 and 4.10 below). Recall the definition of the index set \mathbb{I} in (1.38).

Proposition 4.8 (Proof of Theorem 1.15). *Let $h > 0$, and take let $(m, s) \in \mathbb{I}$ be such that*

$$\text{eq:h-between-hs} \quad (4.60) \quad h^{(m,s)} \leq h < h^{(m,s-1)}.$$

If s is odd, then for N odd

$$\text{eq:max-odd} \quad (4.61) \quad \max_{1 \leq i \leq N^n} \mathcal{H}(h; \gamma_i) = \mathcal{H}(h; \gamma_{\lfloor sN^m/2 \rfloor})$$

and for N even

$$\text{eq:max-even,even} \quad (4.62) \quad \max_{1 \leq i \leq N^n} \mathcal{H}(h; \gamma_i) = \mathcal{H}(h; \gamma_r),$$

where r is given in (4.54). If s is even, then for N odd

$$\text{eq:max-odd,even} \quad (4.63) \quad \max_{1 \leq i \leq N^n} \mathcal{H}(h; \gamma_i) = \mathcal{H}(h; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}),$$

and for N even

$$\text{eq:max-even,odd} \quad (4.64) \quad \max_{1 \leq i \leq N^n} \mathcal{H}(h; \gamma_i) = \mathcal{H}(h; \gamma_{r'}),$$

where r' is obtained by replacing s by $s - 1$ in the leading term in (4.54). If $h \geq \tilde{J}(1 - \frac{1}{N})n$, then $\max_{1 \leq i \leq N^n} \mathcal{H}(h; \gamma_i) = \mathcal{H}(h; \gamma_0)$. If the inequality on the left side of h in (4.60) is also strict, then these are the unique maxima.

Proof. We give the proof for N odd and s even, the proof for all other cases being similar. From (4.3) and Proposition 4.6 we have that, for all $i \leq \lfloor (s - 1)N^m/2 \rfloor + 1$,

$$\text{eq:h-gen-max-ineq} \quad (4.65) \quad \mathcal{H}(h; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}) - \mathcal{H}(h; \gamma_i) \geq \mathcal{H}(h^{(m,s-1)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}) - \mathcal{H}(h^{(m,s-1)}; \gamma_i) \geq 0$$

and, for $i \geq \lfloor (s - 1)N^m/2 \rfloor + 1$,

$$\text{eq:h-gen-max-ineq2} \quad (4.66) \quad \mathcal{H}(h; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}) - \mathcal{H}(h; \gamma_i) \geq \mathcal{H}(h^{(m,s)}; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}) - \mathcal{H}(h^{(m,s)}; \gamma_i) \geq 0.$$

This proves the first claim. If the inequalities in (4.60) are both strict, then both (4.65) and (4.66) are strict whenever $i \neq \lfloor (s - 1)N^m/2 \rfloor + 1$. \square

Remark 4.9. It is easy to check that if we take $h = \tilde{J}((1 - \frac{1}{N})(n - m) - (s - 1)\frac{1}{N}) = h^{(m,s)}$ or $h = h^{(m,s-1)} = \tilde{J}((1 - \frac{1}{N})(n - m) - (s - 2)\frac{1}{N})$ in (4.60), then (4.63) and (4.61) remain true.

Proposition 4.10 (Proof of Theorem 1.11). *Let $h > 0$, and let m and s satisfy (4.60).*

(1) *Suppose that N is odd. For s even,*

$$(4.67) \quad \Gamma^* = \frac{\tilde{J}}{4N} \left(N^m \left[2s \left(N - \frac{s}{2} + 1 \right) - N - 1 \right] + N - 2s + 1 \right) + \frac{1}{2} \left(\tilde{J} \left(1 - \frac{1}{N} \right) (n - m - 1) - h \right) (N^m (s - 1) + 1)$$

while for s odd

$$(4.68) \quad \Gamma^* = \frac{\tilde{J}}{4N} \left(N^m \left[2s \left(N - \frac{s}{2} \right) + N \right] + N - 2s - 1 \right) + \frac{1}{2} \left(\tilde{J} \left(1 - \frac{1}{N} \right) (n - m - 1) - h \right) (sN^m + 1).$$

(2) *Suppose that N is even. For s odd,*

$$(4.69) \quad \begin{aligned} \Gamma^* &= \Gamma_s^* \\ &= \frac{\tilde{J}}{2} N^{1+Q(m)} \left(\frac{N^{m-2+Q(m)} - 1}{N^2 - 1} \right) + \tilde{J} \left(\frac{1}{2} \left(\frac{N^m - 1}{N - 1} \right) - N^{Q(m)} \left(\frac{N^{m-Q(m)} - 1}{N^2 - 1} \right) \right) \\ &\quad \times (N - s) \\ &+ \left[\frac{N}{4} \left(\frac{N^m - 1}{N - 1} \right) - N^{Q(m)} \left(\frac{N^{m-Q(m)} - 1}{N^2 - 1} \right) + N^{m-1} \left(\frac{s-1}{2} \right) \left(N - \frac{s-1}{2} \right) \right] \\ &+ \left(\left(\frac{s-1}{2} \right) N^m + \frac{N}{2} \left(\frac{N^m - 1}{N - 1} \right) - N^{1+Q(m)} \left(\frac{N^{m-Q(m)} - 1}{N^2 - 1} \right) \right) \\ &\quad \times \left(\tilde{J} \left(1 - \frac{1}{N} \right) (n - m - 1) - h \right), \end{aligned}$$

while for s even,

$$\text{eq:shifted-Gamma} \quad (4.70) \quad \begin{aligned} \Gamma^* &= \Gamma_{s-1}^* + \left(h^{(s-1)} - h \right) \\ &\quad \times \left(sN^m - \left(\frac{s-1}{2} \right) N^m - \left(\frac{N}{2} \right) \left(\frac{N^m - 1}{N - 1} \right) + N^{1+Q(m)} \left(\frac{N^{m-Q(m)} - 1}{N^2 - 1} \right) \right). \end{aligned}$$

Proof. From Proposition 4.8 we have that, for N odd and s even,

$$\text{eq:GammastarNoddseven} \quad (4.71) \quad \Gamma^* = \mathcal{H}(h; \gamma_{\lfloor (s-1)N^m/2 \rfloor + 1}) - \mathcal{H}(h; \square),$$

where we also note that

$$(4.72) \quad \lfloor (s-1)N^m/2 \rfloor + 1 = \lfloor (s-1)/2 \rfloor N^m + 1 + \sum_{i=0}^{m-1} \left\lfloor \frac{N}{2} \right\rfloor N^i = \left(\frac{s}{2} - 1\right) N^m + \frac{1}{2}(N^m + 1).$$

We can now use this decomposition together with (4.22) to calculate Γ^* (after a fair deal of tedious computations). For the case where N is odd and s is odd, Γ_s^* is calculated in the same manner, while (4.70) follows immediately from (4.3). \square

4.4. Proof of Theorem 1.17. In this section we identify the configurations in the sets U_σ^- and U_σ^+ defined in Lemma 1.6 and compute the prefactor K^* (Corollary 4.12 and Proposition 4.13 below).

Let M be the volume of the critical configurations, whose value was determined in Proposition 4.8 (i.e., $M = \lfloor sN^{\hat{m}}/2 \rfloor$ if N is odd and s is odd, etc.). Recall that v_M is the last vertex flipped (from -1 to $+1$) in obtaining the configuration γ_M along the path γ . Let $b \geq 1$ and let w be any vertex such that $d(w, v_M) = b$. Define the configuration σ_b by

$$(4.73) \quad \sigma_b(v) = \begin{cases} \gamma_M(v), & v \neq w, \\ -\gamma_M(v), & v = w. \end{cases}$$

Assuming that h satisfies (4.60) with strict inequalities, we know from Proposition 4.8 that any uniformly optimal path attains a unique global maximum. Hence if $b = 1$, then $\mathcal{H}(\sigma_b) < \mathcal{H}(\gamma_M)$, since $\mathcal{H}(\sigma_b) \in \{\mathcal{H}(\gamma_{M-1}), \mathcal{H}(\gamma_{M+1})\}$. The following lemma shows that if $N \neq 2, 4$ and $m \geq 1$, then the only neighbours of γ_M with lower energy are those obtained by flipping a vertex at distance $b = 1$.

Lemma 4.11. *Let σ_b be defined as in (4.73). Suppose that $N \neq 2, 4$ and $m = \hat{m} \geq 1$. Then $\mathcal{H}(\sigma_b) > \mathcal{H}(\gamma_M)$ whenever $b \geq 2$.*

Proof. We first consider $\sigma_b(w) = -1$, where w is the vertex at which σ_b differs from γ_M . Note that $b \leq m + 1$, since there are no $+1$ -valued vertices in γ_M that are at distance larger than $m + 1$ from each other. By (3.36) we have that

$$(4.74) \quad \begin{aligned} \mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_M) &= \tilde{J}(b-1) \left(1 - \frac{1}{N}\right) + \tilde{J}N^{-b} \left(2 \sum_{i=0}^{b-1} a_i N^i - N^b - N^{b-1}\right) \\ &\quad + \frac{\tilde{J}}{N} \sum_{i=b}^{n-1} (2a_i - N + 1) + h. \end{aligned}$$

If $b = m + 1$, then the right-hand side gives

$$(4.75) \quad \begin{aligned} &\tilde{J} \left((b-1) \left(1 - \frac{1}{N}\right) + N^{-b} \left(2 \sum_{i=0}^{b-1} a_i N^i - N^b - N^{b-1}\right) - \left(1 - \frac{1}{N}\right) (n - m - 1) + \frac{h}{\tilde{J}} \right) \\ &\geq \tilde{J} \left((m+1) \left(1 - \frac{1}{N}\right) + N^{-m-1} (2M - N^{m+1} - N^m) - (s-1) \frac{1}{N} \right), \end{aligned}$$

where the inequality follows from the bounds on h in (4.60). It is easy to see from the value of M in Theorem 1.15 that the above is strictly larger than

$$(4.76) \quad \begin{aligned} &\tilde{J} \left((m+1) \left(1 - \frac{1}{N}\right) + \frac{1}{N} ((s-1 - Q(s+1)) - N - 1) - (s-1) \frac{1}{N} \right) \\ &\geq \tilde{J} \left(m \left(1 - \frac{1}{N}\right) - \frac{2}{N} \right) \geq 0. \end{aligned}$$

Hence we conclude that for $b = m + 1$ and $\sigma_b(w) = -1$ the claim of the lemma holds.

Now assume that $b \leq m$. If N is odd, then $a_0 = \frac{N-1}{2} + Q(s+1)$ and $a_i = \frac{N-1}{2}$ for $1 \leq i \leq m-1$, while $a_m = \lfloor \frac{s-1}{2} \rfloor = \frac{s-1-Q(s+1)}{2}$ and $a_i = 0$ for $i > m$. If h satisfies (4.60) for some $1 \leq s \leq N-1$

and $2 \leq m \leq n-1$ (we do not need to consider the case $m=1$ because $m \geq b \geq 2$), then this implies

$$(4.77) \quad \begin{aligned} \mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_M) &= \tilde{J} \left((b-n+m) \left(1 - \frac{1}{N} \right) + N^{-b} (2Q(s+1) - 1) \right. \\ &\quad \left. - \frac{1}{N} (N-s+1+Q(s+1)) + \frac{h}{\tilde{J}} \right) \end{aligned}$$

and hence $\mathcal{H}(\sigma_b) \leq \mathcal{H}(\gamma_k)$ if and only if

$$(4.78) \quad \begin{aligned} b &\leq 1 + \left(1 - \frac{1}{N} \right)^{-1} \left(\frac{1}{N} (N-s+1+Q(s+1)) \right. \\ &\quad \left. + \left(1 - \frac{1}{N} \right) (n-m-1) - N^{-b} (2Q(s+1) - 1) - \frac{h}{\tilde{J}} \right). \end{aligned}$$

From (4.60) we have that the right-hand side of (4.78) is less than or equal to

$$(4.79) \quad \frac{Q(s+1)+1}{N} - N^{-b} (2Q(s+1) - 1)$$

and hence is less than 2 when $N \geq 3$. This implies that $\mathcal{H}(\sigma_b) > \mathcal{H}(\gamma_k)$ when $b \geq 2$. If N is even, then

$$(4.80) \quad \begin{aligned} (\mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_M)) \tilde{J}^{-1} &= (b-1) \left(1 - \frac{1}{N} \right) + N^{-b} \left(2 \sum_{i=0}^{b-1} a_i N^i - N^b - N^{b-1} \right) \\ &\quad - \frac{1}{N} (N-s+1+Q(s+1)) - \frac{1}{N} (N-1) (n-m-1) \\ &\quad + \frac{1}{N} (1 - Q(s+m) - Q(s+b+1)) + \frac{h}{\tilde{J}}, \end{aligned}$$

and hence $\mathcal{H}(\sigma_b) \leq \mathcal{H}(\gamma_k)$ if and only if

$$(4.81) \quad \begin{aligned} (b-1) \left(1 - \frac{1}{N} \right) &\leq 1 - \frac{s}{N} + \frac{Q(s+1)}{N} + \left(1 - \frac{1}{N} \right) (n-m-1) \\ &\quad + \frac{Q(s+m)}{N} + \frac{Q(s+b+1)}{N} - \frac{h}{\tilde{J}} - N^{-b} \left(2 \sum_{i=0}^{b-1} a_i N^i - N^b - N^{b-1} \right). \end{aligned}$$

Since h satisfies (4.60), this implies

$$(4.82) \quad \begin{aligned} b &\leq 1 + \left(1 - \frac{1}{N} \right)^{-1} \left(\frac{Q(s+1) + Q(s+b+1) + Q(s+m)}{N} \right. \\ &\quad \left. - N^{-b} \left(2 \sum_{i=0}^{b-1} a_i N^i - N^b - N^{b-1} \right) \right) \\ &\leq 1 + \left(1 - \frac{1}{N} \right)^{-1} \left(\frac{Q(s+1) + Q(s+b+1) + Q(s+m) + 2Q(s+m-b)}{N} - R_b \right), \end{aligned}$$

where $R_b = N^{-b} \left(\frac{1}{N-1} (N^{b-1} - N - 2N^{b-2}) \right)$. The right-hand side is less than 2 when $N \geq 6$, in which case $\mathcal{H}(\sigma_b) > \mathcal{H}(\gamma_k)$ when $b \geq 2$.

Now suppose that $\sigma_b(w) = +1$. Let us first consider the case when N is odd. Suppose that $b > m$. Then by (3.38)

$$\begin{aligned}
(4.83) \quad & \mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_k) \\
&= \tilde{J} \left(1 - \frac{1}{N}\right) (b-1) + \tilde{J} N^{b-1} \left(N^b - 2 \sum_{i=0}^{b-1} a_i N^i - N^{b-1}\right) + \sum_{i=b}^{n-1} \tilde{J}_{i+1} N^i (N - 2a_i - 1) - h \\
&= \tilde{J} \left(\left(1 - \frac{1}{N}\right) (b-1) + N^{b-1} (N^b - (s - Q(s+1)) N^m + 1 - 2Q(s+1) - N^{b-1}) \right. \\
&\quad \left. + \left(1 - \frac{1}{N}\right) (n-b) - \frac{h}{\tilde{J}} \right).
\end{aligned}$$

From (4.60) it follows that this is larger than or equal to

$$\begin{aligned}
(4.84) \quad & \tilde{J} \left(N^{-b} (N^b - (s - Q(s+1)) N^m + 1 - 2Q(s+1) - N^{b-1}) \right. \\
&\quad \left. + \left(1 - \frac{1}{N}\right) (m-1) + (s-2) \frac{1}{N} \right) > 0.
\end{aligned}$$

Hence, the inequality $\mathcal{H}(\sigma_b) \leq \mathcal{H}(\gamma_k)$ is at most possible for $b \leq m$. In this case we get that $\mathcal{H}(\sigma_b) \leq cH(\gamma_k)$ if and only if

$$\begin{aligned}
\text{eq:bineq+Nodd} \quad (4.85) \quad & b \leq 1 + \left(1 - \frac{1}{N}\right)^{-1} \left(\frac{h}{\tilde{J}} - \left(1 - \frac{1}{N}\right) (n-m-1) \right. \\
&\quad \left. - \frac{1}{N} (N - s + Q(s+1) - 1) - N^{-b} (1 - 2Q(s+1)) \right).
\end{aligned}$$

Once again, from (4.60) it follows that (4.85) is satisfied if and only if

$$(4.86) \quad b \leq 1 + \left(1 - \frac{1}{N}\right)^{-1} \left(-\frac{1}{N} (Q(s+1) - 1) - N^{-b} (1 - 2Q(s+1)) \right) < 2 \quad \forall N \geq 2.$$

Similarly, if N is even, then for $b > m$ we get

$$\begin{aligned}
(4.87) \quad & \mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_M) \geq \tilde{J} \left(\left(1 - \frac{1}{N}\right) (b-1) - \frac{1}{N(N-1)} + \frac{Q(s+1)}{N} + \frac{Q(s+m)}{N} \right. \\
&\quad \left. + \frac{Q(s+b+1)}{N} + \frac{Q(s+m-b)}{N} - \frac{2}{N} \right),
\end{aligned}$$

which is larger than 0 when $N \geq 4$. Thus, once again we only need to consider $b \leq m$, for which $\mathcal{H}(\sigma_b) - \mathcal{H}(\gamma_k) \leq 0$ if and only if

$$\begin{aligned}
\text{eq:bineq+Neven} \quad (4.88) \quad & b \leq 1 + \left(1 - \frac{1}{N}\right)^{-1} \left(-\frac{1}{N} (N - s + 1 + Q(s+1)) - \left(1 - \frac{1}{N}\right) (n-m-1) \right. \\
&\quad \left. + \frac{1}{N} (1 - Q(s+m) - Q(s+b+1)) + \frac{h}{\tilde{J}} - N^{-b} \left(N^b - 2 \sum_{i=0}^{b-1} a_i N^i - N^{b-1} \right) \right) \\
&\leq 1 + \left(1 - \frac{1}{N}\right)^{-1} \left(\frac{1}{N} (1 - Q(s+m) - Q(s+b+1) - Q(s+1)) \right. \\
&\quad \left. - 2Q(s+m-b) \right) + \frac{1}{N(N-1)},
\end{aligned}$$

which is less than 2 when $N \geq 4$. □

The prefactor K^* can now be easily computed.

Corollary 4.12. *Suppose that $N \neq 2, 4$ and $m \geq 1$. Then*

$$\stackrel{\text{eq:corollary 1/K}}{(4.89)} \quad \frac{1}{K^*} = a_0 N^{n-\hat{m}-2} \prod_{i=0}^{\hat{m}} \binom{N}{a_i} (N - a_i).$$

Proof. By Lemma 4.11 we have that, for all $\sigma \in \mathcal{C}^*$,

$$(4.90) \quad \begin{aligned} |U_\sigma^-| &= |w \in \Lambda_N^n: d(w, v_M) = 1, \gamma_M(w) = -1| = a_0, \\ |U_\sigma^-| &= |w \in \Lambda_N^n: d(w, v_M) = 1, \gamma_M(w) = -1| = N - a_0. \end{aligned}$$

Furthermore, it is a simple combinatorial fact that

$$(4.91) \quad \begin{aligned} |\mathcal{C}^*| &= |\{\sigma \in \Omega: \sigma = \varphi(\gamma_M) \text{ for some isometric bijection } \varphi: \Lambda_N^n \rightarrow \Lambda_N^n\}| \\ &= N^{n-\hat{m}-1} (N - a_0)^{-1} \prod_{i=0}^{\hat{m}} \binom{N}{a_i} (N - a_i). \end{aligned}$$

Equation (4.89) now follows from Lemmas 1.6 and 3.6. □

We can also investigate what the prefactor amounts to when the precondition of Corollary 4.12 is not satisfied. For this, we define

$$(4.92) \quad \begin{aligned} O_d &= \{1\} \cup \{2 \leq b \leq \hat{m}: b \text{ satisfies inequality (4.78)}\}, \\ O_u &= \{1\} \cup \{2 \leq b \leq \hat{m}: b \text{ satisfies inequality (4.85)}\}, \end{aligned}$$

and

$$(4.93) \quad \begin{aligned} E_d &= \{1\} \cup \{2 \leq b \leq \hat{m}: b \text{ satisfies inequality (4.81)}\}, \\ E_u &= \{1\} \cup \{2 \leq b \leq \hat{m}: b \text{ satisfies inequality (4.88)}\}. \end{aligned}$$

Then the following is immediate from Lemmas 1.6 and 3.6.

Proposition 4.13. *Suppose that h satisfies*

$$h^{(m,s)} < h < h^{(m,s-1)}$$

for some $(m, s) \in \mathbb{I}$. If N is odd, then the prefactor K^* is given by

$$\stackrel{\text{eq:1/Kstar}}{(4.94)} \quad \frac{1}{K^*} = \frac{[\sum_{i \in O_d} a_{i-1} N^{i-1}] [\sum_{i \in O_u} (N^i - a_{i-1} N^{i-1})]}{[\sum_{i \in O_d} a_{i-1} N^{i-1}] + [\sum_{i \in O_u} (N^i - a_{i-1} N^{i-1})]} \frac{N^{n-\hat{m}-1}}{N - a_0} \prod_{i=0}^{\hat{m}} \binom{N}{a_i} (N - a_i),$$

where $a_0 = \frac{N-1}{2} + 1$, $a_i = \frac{N-1}{2}$ for $i = 1, \dots, \hat{m} - 1$ and $a_{\hat{m}} = \frac{s-1-(s+1) \bmod 2}{2}$. If N is even, then the summations in (4.94) are over E_d and E_u , respectively, and the terms a_i are replaced by \bar{a}_i defined in (4.54).

REFERENCES

- [1] A. Bovier and F. den Hollander, *Metastability – A Potential-Theoretic Approach*, Grundlehren der mathematischen Wissenschaften 351, Springer, 2015.
- [2] S. Dommers, Metastability of the Ising model on random regular graphs at zero temperature, to appear in Probab. Theory Relat. Fields.
- [3] S. Dommers, F. den Hollander, O. Jovanovski and F.R. Nardi, Metastability for Glauber dynamics on random graphs, to appear in Ann. Appl. Probab.
- [4] O. Jovanovski, Metastability for the Ising Model on the hypercube, to appear in J. Stat. Phys.
- [5] E. Olivieri and M.E. Vares, *Large Deviations and Metastability*, Encyclopedia of Mathematics and its Applications 100, Cambridge University Press, Cambridge, 2005.